

FOR DISTRIBUTION AND USE ONLY IN THE STATES OF IA, IL, IN, KS, KY, MI,
NE (fields south of Route 30 and east of route 281), OH and
WI (only in the counties of Grant, Lafayette, Green, Rock, Walworth, Kenosha and Racine)

ENCOMPASS HERBICIDE

PLUS CANOPY® EX HERBICIDE PLUS 2,4-D HERBICIDE TANK MIX FOR FALL APPLICATION OR
UP TO 45 DAYS PRIOR TO SOYBEAN PLANTING FOR WEED CONTROL IN SOYBEANS

EPA Reg. Nos. 59639-99-55467 & 352-635
FIFRA SECTION 2(ee) RECOMMENDATION

DIRECTIONS FOR USE

Herbicide	Rate
<i>Encompass</i> Herbicide	1.5 to 2 oz./A
plus	plus
Canopy EX	1.1 oz./A
plus	plus
2,4-D ester or 2,4-D amine	0.5 to 1.0 lb. a.i./A

Adjuvant: Crop oil concentrate at 1 pt./A or NIS at 0.25% v/v.

NOTE: COC is the preferred adjuvant. NIS should only be used when recommended by local Tenkoz personnel.

Weeds Controlled by *Encompass plus Canopy EX plus 2,4-D*

Common Name	Scientific Name	Postemergence/Burndown	Residual
Chickweed		Weeds 3 inches or less	
Common	<i>Stellaria media</i>	Yes	Yes
Mouseear	<i>Cerastium vulgatum</i>	Yes	Yes
Dandelion	<i>Taraxacum officinale</i>	Yes ¹	Yes
Henbit	<i>Lamium amplexicaule</i>	Yes	Yes
Marestail/Horseweed	<i>Conyza Canadensis</i>	Yes	Yes
Groundsel, Cressleaf	<i>Senecio glabellus</i>	-	Yes
Purple Deadnettle	<i>Lamium purpureum</i>	Yes	Yes
Annual Fleabane	<i>Erigeron annuus</i>	Yes	-
		Weeds 12 inches or less	
Carolina Geranium	<i>Geranium carolinianum</i>	Yes	-
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>	Yes	Yes
Mustard, Wild	<i>Brassica Kaber</i>	Yes	Yes
Shepherd's-purse	<i>Capsella bursa-pastoris</i>	Yes	Yes

(1) 1 lb. a.i./A of 2,4-D or a tank mix of 0.5 lb. a.i./A of 2,4-D plus 0.2 lb. a.i./A of dicamba should be used for control of emerged dandelion. **NOTE:** Only use the 2,4-D plus dicamba tank mix for fall-applied applications.

**GENERAL USE RESTRICTIONS AND LIMITATIONS FOR *ENCOMPASS* HERBICIDE PLUS CANOPY EX
HERBICIDE PLUS 2,4-D HERBICIDE TANK MIX FOR FALL-APPLIED WEED CONTROL**

1. Do not apply to frozen or snow-covered soil.
2. Do not perform any tillage operation after application or residual weed control will be reduced.
3. Refer to the Canopy EX, 2,4-D, and dicamba labels for additional weeds controlled and rotational restrictions.
4. Application must be made no earlier than October 15 or when soil temperature falls below 50°F at a two-inch depth to maintain residual weed control into the spring (May 1) or up until planting, whichever comes first.
5. Abnormally warm or wet winters will reduce the length of weed control observed in the spring.
6. Only soybeans may be planted in the spring following applications made during the previous year.

Follow all applicable directions, restrictions and precautions on the EPA-registered labels.

It is a violation of Federal Law to use these products in a manner inconsistent with its labeling.

THIS RECOMMENDATION EXPIRES DECEMBER 31, 2006

Encompass -- TM of Tenkoz Inc.
Canopy EX -- Reg. TM of DuPont Company