



Avalanche™ Bulk Pak Herbicide

**For Agricultural or Commercial Use Only
NOT FOR SALE OR USE IN CALIFORNIA**

EPA Reg. No. 279-3242-1381 EPA Est. 279-IL-1

Active Ingredient: By Wt.

Carfentrazone-ethyl: Ethyl α ,2-dichloro-5-[4-(difluoromethyl)-4,5-dihydro-3-methyl-5-oxo-1H-1,2,4-triazol-1-yl]-4-fluorobenzenepropanoate 21.3%

Inert Ingredients: 78.7%
100.0%

This product contains 1.9 pounds active ingredient per gallon.

Contains Petroleum Distillates

U.S. Patent No. 5,125,958

**KEEP OUT OF REACH OF CHILDREN
CAUTION**

FIRST AID

If Inhaled: Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth, if possible. Call a poison control center or doctor for further treatment advice.

If on Skin or Clothing: Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.

If in Eyes: Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

If Swallowed: Call a poison control center or doctor immediately for treatment advice. Have person sip a glass of water if able to swallow. Do not induce vomiting unless told to do so by the poison control center or doctor. Do not give anything by mouth to an unconscious person.

HOTLINE NUMBER

Have the product container or label with you when calling a poison control center or doctor, or going for treatment. You may also contact 1-800-331-3148 for emergency medical treatment information.

Distributed by:

Agriance, LLC

P.O. Box 64089

St. Paul, MN 55164-0089

3/8/2002

Note to Physician: Carfentrazone-ethyl 2EW is expected to have low oral and dermal toxicity, and moderate inhalation toxicity. It is expected to be slightly irritating to the skin and minimally irritating to the eyes. Treatment is otherwise controlled removal of exposure followed by symptomatic and supportive care. This product may pose an aspiration pneumonia hazard.

See other panels for additional precautionary information.

PRECAUTIONARY STATEMENTS

Hazards to Humans (and Domestic Animals)

Caution

Harmful if swallowed, absorbed through the skin or inhaled. Causes moderate eye irritation. Avoid breathing dust. Avoid contact with eyes, skin or clothing. Wash thoroughly with soap and water after handling.

Personal Protective Equipment (PPE)

Applicators and other handlers must wear: long-sleeved shirt and long pants, waterproof gloves, and shoes plus socks.

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product. Do not reuse them. Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

User Safety Recommendations:

Users should:

- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

Environmental Hazards

Carfentrazone-ethyl is very toxic to algae and moderately toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the high water mark, except as specified on this label. Do not contaminate water when disposing of equipment washwaters.

Physical/Chemical Hazards

FLAMMABLE Do not use or store near heat or open flame.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is: Coveralls, chemical resistant gloves, such as polyethylene or polyvinyl chloride, and shoes plus socks.

STORAGE AND DISPOSAL

Pesticide Storage

Not for use or storage in or around the house.

Keep out of reach of children and animals. Store in original containers only. Store in a cool, dry place and avoid excess heat. Carefully open containers. After partial use, replace lids and close tightly. Do not put granule or dilute material into food or drink containers. Do not contaminate other pesticides, fertilizers, water, food, or feed by inappropriate storage or disposal.

In case of spill, avoid contact, isolate area and keep out unprotected persons and animals. Confine spills. Call FMC: (800) 331-3148.

To confine spill: Dike surrounding area, sweep up spillage. Dispose of in accordance with information given under Pesticide Disposal. Wash spill area with water, absorb with sand, cat litter or commercial clay, sweep up and dispose of in an approved manner. Place damaged container in a larger holding container. Identify contents per required hazardous waste labeling regulations.

Pesticide Disposal

Waste resulting from the use of this product may be disposed of at an approved waste disposal facility.

Container Disposal

Triple rinse (or equivalent). Then offer for approved pesticide container recycling program, or puncture and dispose of in an approved waste disposal facility. Provided on site incineration is allowed by state and local authorities, stay out of smoke.

GENERAL INFORMATION

Avalanche Bulk Pak is a water emulsion formulation. It is designed to be mixed with water and applied to corn (field, sweet, seed, popcorn, and silage), grain sorghum, soybeans, rice, wheat, barley, and oats for selective postemergence control of broadleaf weeds. Weed control is best when the product is applied to actively growing weeds up to 4 inches in height. Avalanche Bulk Pak is a contact herbicide with little or no residual activity at recommended use rates.

Avalanche Bulk Pak is rapidly absorbed through the foliage of plants. To avoid significant crop response, applications should not be made within 6 - 8 hours of either rain or irrigation. Within a few hours following application, the foliage of susceptible weeds show signs of desiccation, and in subsequent days necrosis and death of the plant. Due to environmental conditions and with certain spray tank additives, some herbicidal symptoms may appear on the crop. However, the crop recovers quickly with no loss in yield.

Extremes in environmental conditions e.g. temperature and moisture, soil conditions, and cultural practices may affect the activity of Avalanche Bulk Pak. Under warm moist conditions, herbicide symptoms may be accelerated. While under very dry conditions, the expression of herbicide symptoms is delayed, and weeds hardened off by drought are less susceptible to Avalanche Bulk Pak.

Tank Mixtures

Avalanche Bulk Pak may be tank mixed with other herbicides to control weeds not listed on this label. Read and follow all manufacturer's label recommendations for the companion herbicide except for specific recommendations on this label. Tank mixtures of Avalanche Bulk Pak with EC formulations of other crop protection products, crop oil concentrate, methylated seed oil, silicone based adjuvants, 28% nitrogen or ammonium sulfate may increase leaf speckling.

With adjuvants

Use a non-ionic surfactant (NIS) having at least 80% active ingredient strength at 0.25% v/v (2 pints per 100 gallons of spray solution) OR a non-ionic surfactant at 0.25% v/v (2 pints per 100 gallons) plus 28% nitrogen (UAN) at 2 to 4 quarts per 100 gallons or ammonium sulfate (AMS) at 2-4 pounds per acre where recommended by those companion herbicides listed on this label. In the latter case, the level of leaf speckling may be higher than with NIS alone. Crop oil or crop oil plus either 28% nitrogen or ammonium sulfate may be used with companion herbicides listed on this label and may be recommended under very dry conditions.

Mixing and Loading Instructions:

Fill the spray tank 3/4 full with clean water. Make sure the agitation

system is operating. Add the recommended amount of Avalanche Bulk Pak and complete filling the spray tank to the desired level. The spray tank agitation should be sufficient to ensure uniform spray mixture during application. When tank mixing with other products, Avalanche Bulk Pak should be mixed first in the spray tank. After the product is thoroughly mixed, add the other products as specified on their label. Ensure the compatibility of other products with Avalanche Bulk Pak before mixing them together in the spray tank.

Do not use with tank additives that alter the pH of the spray solution below pH 5 or above pH 8. Buffer spray solution to alter the pH range as appropriate.

Product Application Guidelines:

Thorough weed coverage with the spray mixture is essential for optimum weed control. Do not apply when conditions are conducive to spray drift, poor spray deposition or poor weed coverage.

When Avalanche Bulk Pak is applied as a tank mixture with other herbicides, read and follow the label directions of the other product if they are more restrictive than those of the Avalanche Bulk Pak label.

Ground Applications:

Utilize a boom and nozzle sprayer equipped with the appropriate nozzles, spray tips and screens and adjusted to provide optimum spray distribution and coverage at the appropriate operating pressures. Utilize nozzles which produce minimal amounts of fine spray droplets. Do not exceed 30 psi spray pressure unless otherwise required by the manufacturer of drift reducing nozzles. Apply a minimum of 10 gallons of finished spray per acre. Higher spray volumes are required when there is a dense weed population or crop canopy. Sprayers should be adjusted to position spray tips a minimum of 18 inches above the crop and operated to avoid the application of excessive herbicide rates directly over the rows and/or into the whorl of treated crop plants. Be aware that overlaps and slower ground speeds while starting, stopping or turning while spraying may result in higher application rates and possible crop response.

Aerial Application:

Use nozzle types and arrangements which will provide optimum coverage while producing a minimal amount of fine droplets. Apply at a minimum of 3 gallons of finished spray per acre. Higher spray volumes are required when there is a dense weed population or crop canopy.

Spray Equipment Clean-Out:

After spraying Avalanche Bulk Pak and before using sprayer equipment for any other applications, the sprayer must be thoroughly cleaned using the following procedure. In addition, users must take appropriate steps to ensure proper equipment clean-out for any other products mixed with Avalanche Bulk Pak as explained on the other product labels.

1. Drain sprayer tank, hoses, and spray boom and thoroughly rinse with clean water the inside of the sprayer tank to remove sediment and residues, and thoroughly flush sprayer hoses, boom and nozzles with clean water.
2. Fill the tank 1/2 full with clean water, and add appropriate detergent (follow manufacturer's directions for use). Fill the tank to capacity and operate the sprayer for 15 minutes to flush hoses, boom and nozzles.
3. Drain the sprayer system. Rinse the tank with clean water and flush through the hoses, boom, and nozzles. Remove and clean spray tips and screens separately.
4. Properly dispose of all cleaning solution and rinsate in accordance with Federal, State, and local regulations and guidelines.

Aerial Spray Drift Management

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR AND THE GROWER.

The interaction of many equipment-and-weather-related factors determine the potential for spray drift. The applicator and the grower are responsible for considering all these factors when making decisions.

The following drift management requirements must be followed to avoid off-target movement from aerial applications to agricultural field crops. These requirements do not apply to forestry applications, public health uses or to applications of dry materials.

1. The distance of the outer most nozzles on the boom must not exceed 3/4 the length of the wingspan or rotor.

2. Nozzles must always point backward parallel with the air stream and never be pointed downwards more than 45 degrees.

Where states have more stringent regulations, they must be observed. The applicator should be familiar with and take into account the information covered in the Aerial Drift Reduction Advisory.

INFORMATION ON DROPLET SIZE

The most effective way to reduce drift potential is to apply large droplets. The best drift management strategy is to apply the largest droplets that provide sufficient coverage and control. Applying larger droplets reduces drift potential, but will not prevent drift if applications are made improperly, or under unfavorable environmental conditions (See Wind, Temperature and Humidity, and Temperature Inversions).

CONTROLLING DROPLET SIZE

Volume - Use high flow rate nozzles to apply the highest practical spray volume. Nozzles with higher rated flows produce larger droplets.

Pressure - Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

Number of nozzles - Use the minimum number of nozzles that provide uniform coverage.

Nozzle Orientation - Orienting nozzles so that the spray is released parallel to the airstream produces larger droplets than other orientations and is the recommended practice. Significant deflection from horizontal will reduce droplet size and increase drift potential.

Nozzle Type - Use a nozzle type that is designed for the intended application. With most nozzle types, narrower spray angles produce larger droplets. Consider using low drift nozzles. Solid stream nozzles oriented straight back produce the largest droplets and the lowest drift.

BOOM LENGTH

For some use patterns, reducing the effective boom length to less than 3/4 of the wingspan or rotor length may further reduce drift without reducing swath width.

APPLICATION HEIGHT

Applications should not be made at a height greater than 10 feet above the top of the target plants unless a greater height is required for aircraft safety. Making applications at the lowest height that is safe reduces exposure of droplets to evaporation and wind.

SWATH ADJUSTMENT

When applications are made with a crosswind, the swath will be displaced downwind. Therefore, on the up and downwind edges of the field, the applicator must compensate for this displacement by adjusting the path of the aircraft upwind. Swath adjustment distance should increase, with increasing drift potential (higher wind, smaller drops, etc.)

WIND

Drift potential is lowest between winds speeds of 3-10 mph. However, many factors, including droplet size and equipment type determine drift potential at any given speed. Application should be avoided below 3 mph due to variable wind direction and high inversion potential. NOTE: Local terrain can influence wind patterns. Every applicator should be familiar with local wind patterns and how they affect spray drift.

TEMPERATURE AND HUMIDITY

When making applications in low relative humidity, set up equipment to produce larger droplets to compensate for evaporation. Droplet evaporation is most severe when conditions are both hot and dry.

TEMPERATURE INVERSIONS

Applications should not occur during a temperature inversion because drift potential is high. Temperature inversions restrict vertical air mixing, which causes small suspended droplets to remain in a concentrated cloud. This cloud can move in unpredictable directions due to the light variable winds common during inversions. Temperature inversions are characterized by increasing temperatures with altitude and are common on nights with limited cloud cover and light to no wind. They begin to form as the sun sets and often continue into the

morning. Their presence can be indicated by ground fog; however, if fog is not present, inversions can also be identified by the movement of smoke from a ground source or an aircraft smoke generator. Smoke that layers and moves laterally in a concentrated cloud (under low wind conditions) indicates an inversion, while smoke that moves upward and rapidly dissipates indicates good vertical air mixing.

SENSITIVE AREAS

The pesticide should only be applied when the wind is blowing away from adjacent sensitive areas (e.g. residential areas, bodies of water, known habitat for threatened or endangered species, non-target crops).

Post Fallow or Preplant Burndown

Timing and Method of Application

Apply Avalanche Bulk Pak alone or with other herbicides in the post fallow period prior to planting or emergence of labeled crops to control or suppress annual broadleaf weeds. For all products used in tank mixes, refer to the specific product labels for all restrictions on tank mixing and observe all label precautions, instructions and rotational cropping restrictions. Do not apply more than 2 fl oz (0.031 pound active ingredient) per acre per season including post fallow/preplant burndown and labeled crop applications to corn. Do not apply more than 10 fl oz (0.15 pound active ingredient) per acre per season post fallow/preplant burndown and labeled crop applications to sorghum.

For best performance, make applications to actively growing weeds up to 4 inches high or rosettes less than 3 inches across. Coverage is essential for good control. Optimum broad spectrum control of annual and perennial weeds requires a tank mix of a broad spectrum burndown herbicide such as RoundUp Ultra®, Touchdown® or Gramoxone® Extra.

Fallow Systems

Use Avalanche Bulk Pak at 1/4 - 2 fl oz (0.004-0.031 pound active ingredient) per acre in fallow systems. A nonionic surfactant or crop oil concentrate may be used to enhance activity of Avalanche Bulk Pak in fallow systems. Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient or a petroleum or vegetable seed based oil concentrate at 1.5 to 2.0 pints per acre.

Crop Rotation for applications in corn

Corn (field, sweet, seed, popcorn, and silage) and grain sorghum may be planted any time following an application of Avalanche Bulk Pak. Follow rotation statements on tank mix products if they are more restrictive.

CORN: FIELD CORN, SEED CORN, POPCORN, CORN SILAGE, AND SWEET CORN FOR PROCESSING, AND FRESH MARKET SWEET CORN:

TIMING AND METHOD OF APPLICATION

Apply Avalanche Bulk Pak alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to corn in all tillage systems from 30 days before planting up to 8 leaf collar growth stage. Do not apply when conditions favoring drift exist or wind is above 10 mph.

For best performance, make application to actively growing weeds up to 4 inches high and rosettes less than 3 inches across. **Coverage is essential for good control.**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Under dry conditions the use of a crop oil concentrate may improve weed control. The use of a crop oil concentrate may increase leaf speckling on the treated corn leaves.

To control weeds not listed on this label, Avalanche Bulk Pak may be tank mixed with other herbicides registered for use in corn. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions, and rotational cropping restrictions.

Sprayers should be adjusted to position spray tips a minimum of 18 inches above the crop and operated to avoid the application of excessive herbicide rates directly over the rows and/or into the whorl of the corn plant. Be aware that overlaps and slower ground speeds while starting, stopping or turning while spraying may result in higher application rates and possible crop response. Avalanche Bulk Pak may be applied with drop nozzles.

Use Rates

Use Avalanche Bulk Pak at 1/2 fl oz (0.008 pound active ingredient) per acre. Applications should be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre or by air at a minimum finished spray volume of 3 gallons of spray per acre. When applied as directed, Avalanche Bulk Pak will control the following weeds:

Weeds controlled

Common Lambsquarters (**up to 3 inches tall**)

Morningglories (**2-3 true leaves**),

Ivyleaf and Pitted

Nightshade, Eastern Black (**up to 4 inches tall**)

Pigweed, Redroot (**up to 4 inches tall**)

Velvetleaf (**up to 18 inches or up to 36 inches with drop nozzles**)

In addition to the above weeds listed as controlled, Avalanche Bulk Pak will suppress or partially control the following weeds:

Weeds suppressed (up to 4 inches tall)

Cocklebur

Common Ragweed

Common (annual) Sunflower

Field Bindweed

Jimsonweed

Kochia

Morningglory, Entireleaf

Palmer Amaranth

Pennsylvania Smartweed

Pigweed, Smooth

Prickly Sida

Prostrate Spurge

Smooth Groundcherry

Trumpet Creeper

Waterhemp (tall, common)

Do not apply more than 2 fl oz (0.031 pound active ingredient) per acre per season including fallow/preplant burndown and labeled crop applications.

Tank Mixtures

Avalanche Bulk Pak may be tank mixed with other herbicides to control weeds not listed on this label. Read and follow all manufacturer's label recommendations for the companion herbicide except for specific recommendations on this label. When tank mixing Avalanche Bulk Pak with other products, be sure the Avalanche Bulk Pak is mixed in the spray tank water first.

For control of additional broadleaf weeds and grasses, Avalanche Bulk Pak may be tank mixed with 2,4-D (amine), Accent®, Accent Gold®, Atrazine, Banvel®, Basis®, Basis Gold®, Beacon®, Clarity™, Distinct®, Exceed®, Hornet®, Liberty®, Lightning®, Marksman®, Northstar™, Permit®, Poast®, Roundup®, Roundup Ultra®, Scorpion® III, Sencor®, Spirit™, Sterling®, Touchdown®, and Tough®.

When tank mixing Avalanche Bulk Pak with Accent, Accent Gold, Atrazine, Basis Gold, Liberty, Poast®, and Roundup Ultra use adjuvants recommended on the tank mix partner label. These may include nonionic surfactant, crop oil concentrate, 28% nitrogen, ammonium sulfate or combinations of these.

Leaf speckling can occur when Avalanche Bulk Pak is used with certain crop protection products and adjuvants. Refer to the Tank Mixtures and Recommended Adjuvants sections under General Information..

Seed Production

For seed production fields, apply Avalanche Bulk Pak using drop nozzles only. Avoid directing spray solution into the whorl.

Seed corn inbreds have generally shown good tolerance to Avalanche Bulk Pak, however all inbreds have not been tested. Broadcast applications may result in spray being concentrated into the whorl of the plant which will increase leaf response. To minimize application into the whorl of the plants, drop nozzles must be used to direct the spray to the targeted weeds.

Sweet Corn Production

Avalanche Bulk Pak herbicide may be applied to sweet corn, however, the user assumes all responsibility for herbicide tolerance with such use. All hybrids/varieties have not been tested for sensitivity to

Avalanche Bulk Pak nor does Agrilience have access to all seed company or food processor data. Therefore, any crop response arising from the use of Avalanche Bulk Pak on sweet corn is the responsibility of the user. Use Avalanche Bulk Pak only under the recommendation of the seed company, food processor, or State Agricultural Extension Service.

With Atrazine

Avalanche Bulk Pak may be tank mixed at a rate of 1/2 fl oz (0.008 pound active ingredient) per acre with Atrazine 4L or Atrazine 90DF to control the following weeds.

Weeds (up to 4 inches tall)

Amaranth, Palmer

Amaranth, Spiny

Buffalobur

Carpetweed

Cocklebur, Common

Croton, Wolly

Devilsclaw

Eveningprimrose, Cutleaf

Hophornbean Copperleaf

Jimsonweed

Lambsquarters, Common

Morningglory, Entireleaf

Morningglory, Ivyleaf

Morningglory, Pitted

Morningglory, Scarlet

Morningglory, Tall

Nightshade, Eastern Black

Nightshade, Silverleaf

Pigweed, Redroot

Pigweed, Smooth

Pigweed, (Triazine resistant)

Purslane, Common

*Ragweed, Common

Sesbania, Hemp

*Smartweed, annual

Spurred Anoda

Velvetleaf

Venice Mallow

Waterhemp, Common

Waterhemp, Tall

*Ragweed, Giant

*Sunflower, Wild

*Suppression or partial control

Refer to the Atrazine labels for additional weed listings and for higher use rates.

For control of giant and common ragweeds, annual smartweeds, and wild sunflower:

Avalanche Bulk Pak plus Atrazine can be tank mixed with 2,4-D (amine), Banvel® or Clarity herbicides. Add 2,4-D (amine) to the tank mix at 0.125 - 0.25 pound active ingredient per acre or Banvel® or Clarity at 3-4 fluid ounces. per acre. Higher rates of Atrazine, Sterling® or Clarity herbicides can be used, but do not exceed the recommended label use rates allowed by these labels. Add a 0.25% v/v nonionic surfactant (2 pints per 100 gallons) to the tank mixture, or under very dry soil moisture conditions, the use of crop oil concentrate (1% v/v or 1 gallon per 100 gallon spray solution) may improve weed control. However, the use of crop oil concentrate may increase leaf injury. Refer to the Tank Mixture section for information on potential leaf injury.

GRAIN SORGHUM

TIMING AND METHOD OF APPLICATION

This product may not be applied to sweet sorghum. Apply Avalanche Bulk Pak alone or as a tank mixture with other herbicides to emerged and actively growing weeds. Apply to sorghum in all tillage systems from 30 days before planting up through the 6 leaf growth stage. Do not apply when conditions favoring drift exist or wind is above 10 mph.

For best performance, make application to actively growing weeds up to 4 inches tall and rosettes less than 3 inches across. **Coverage is essential for good control.**

Use a nonionic surfactant at 0.25% v/v (2 pints per 100 gallons of spray solution) having at least 80% active ingredient. Postemergence broadcast applications of Avalanche Bulk Pak with crop oil concentrate are not recommended or increased crop response may occur.

To control weeds not listed on this label, Avalanche Bulk Pak may be tank mixed with other herbicides registered for use in grain sorghum. Refer to the other product's label for restrictions on tank mixing, and observe all label precautions, instructions, and rotational cropping restrictions.

Sprayer should be adjusted and operated to avoid the application of excessive herbicide rates directly over the row and/or into the whorl of the sorghum plant.

Broadcast applications of Avalanche Bulk Pak to sorghum with wet foliage or application during periods of adverse environmental conditions such as cool, cloudy, wet, or high humidity can cause increased crop response. Drop nozzles are recommended if

applications are to be made under any of these conditions to limit the amount of product that may get onto sorghum leaves and/or into the sorghum whorl.

When applying Avalanche Bulk Pak postemergence to sorghum grown for seed, the use of drop nozzles is recommended.

Use Rates

Use Avalanche Bulk Pak at 1/2 fl oz (0.008 pound active ingredient) per acre. Applications should be made by ground equipment using a minimum finished spray volume of 10 gallons of spray per acre or by air at a minimum finished spray volume of 3 gallons of spray per acre. When applied as directed, Avalanche Bulk Pak will control the following weeds:

Weeds controlled

Common Lambsquarters (**up to 3 inches**)

Morningglories (**2-3 true leaves**),

Ivyleaf and Pitted

Nightshade, Black (**up to 4 inches**)

Pigweed, Redroot (**up to 4 inches**)

Velvetleaf (**up to 18 inches or up to 36 inches with drop nozzles**)

In addition to the above weeds listed as controlled, Avalanche Bulk Pak will suppress or partially control the following weeds:

Weeds suppressed (up to 4 inches)

Cocklebur

Common Ragweed

Common (annual) Sunflower

Field Bindweed

Jimsonweed

Kochia

Morningglory, Entireleaf

Palmer Amaranth

Pennsylvania Smartweed

Pigweed, Smooth

Prickly Sida

Prostrate Spurge

Smooth Groundcherry

Thistle, Russian

Trumpet Creeper

Waterhemp (tall, common)

Do not apply more than 1 fl oz (0.015 pound active ingredient) per acre per season including fallow/preplant burndown and labeled crop applications.

Tank Mixtures

Avalanche Bulk Pak may be tank mixed with other herbicides to control weeds not listed on this label. Read and follow all manufacturer's label recommendations for the companion herbicide except for specific recommendations on this label. When tank mixing Avalanche Bulk Pak with other products, be sure the Avalanche Bulk Pak is mixed in the spray tank water first.

For control of additional broadleaf weeds and grasses, Avalanche Bulk Pak may be tank mixed with 2,4-D (amine), Atrazine, Banvel®, Clarity™, Laddok®, Paramount, Peak®, Permit®, and Sterling®.

Leaf speckling can occur when Avalanche Bulk Pak is used with certain formulations of crop protection products and adjuvants. Refer to the Tank Mixtures and Recommended Adjuvants sections under General Information.

Dealers Should Sell in Original Packages Only.

Terms of Sale or Use: On purchase of this product buyer and user agree to the following conditions:

Warranty: Agrilience makes no warranty, expressed or implied, concerning the use of this product other than indicated on the label. Except as so warranted the product is sold as is. Buyer and user assume all risk of use and/or handling and/or storage of this material when such use and/or handling and/or storage is contrary to label instructions.

Directions and Recommendations: Follow directions carefully. Timing and method of application, weather and crop conditions, mixture with other chemicals not specifically recommended and other influencing factors in the use of this product are beyond the control of the seller and are assumed by the buyer at his own risk.

Use of Product: Agrilience's recommendations for the use of this product are based upon tests believed to be reliable. The use of this product being beyond the control of the manufacturer, no guarantee, expressed or implied is made as to the effects of such or the results to be obtained if not used in accordance with directions or established safe practice.

Damages: Buyer's or user's exclusive remedy for damages for breach of warranty or negligence shall be limited to direct damages not exceeding the purchase price paid and shall not include incidental or consequential damages.

Accent, Accent Gold, Basis, Basis Gold, - trademarks of E.I. DuPont de Nemours and Company

Gramoxone, Touchdown, Beacon, Exceed, Northstar, Peak, Spirit, Tough - trademarks of Syngenta, Inc.

Lightning - trademark of BASF

Banvel, Clarity, Distinct, Marksman - trademarks of BASF Corporation

Curtail, Hornet, Scorpion - trademarks of Dow Agrosciences, LLC

Laddok, Poast - trademarks of BASF Aktiengesellschaft

Liberty - trademark of Hoechst Schering AgrEvo GmbH

Permit - trademark of Nissan Chemical Industries, Inc.

Roundup, Roundup Ultra - trademarks of Monsanto Company

Sencor, Def - trademarks of Bayer Aktiengesellschaft

Sterling, Avalanche - trademarks of Agrilience, LLC