

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF FLORIDA

**RESTRICTED USE PESTICIDE**

Due to toxicity to aquatic invertebrate animals.  
For retail sale to and use only by Certified Applicators, or persons under their direct supervision, and only for those uses covered by the Certified Applicator's certification.



**MICROMITE® 25WS**

INSECT GROWTH REGULATOR  
EPA REG. NO. 400-471

Active Ingredients: (% by weight)  
Diflubenzuron N-[[[(4-Chlorophenyl)amino]carbonyl]-2,6-difluorobenzamide\* ..... 25.0%  
Inert Ingredients ..... 75.0%

**KEEP OUT OF REACH OF CHILDREN  
CAUTION**

**THIS LABEL IS IN ADDITION TO THE STANDARD FEDERAL LABEL PRINTED ON THE CONTAINER  
DIRECTIONS FOR USE**

**PERSONAL PROTECTIVE EQUIPMENT**

**Mixers and Loaders Using Fixed Wing Aircraft Must Wear:** Long-sleeved shirt and long pants, chemical-resistant gloves such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, PVC or viton; shoes plus socks; dust/mist filtering respirator (MSHA/NIOSH approval number prefix TC-21C or a NIOSH approved respirator with any R, P, or HE filter). Follow the manufacturer's instructions for cleaning / maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

**Applicators and Other Handlers Must Wear:** Long-sleeved shirt and long pants; chemical-resistant gloves, such as barrier laminate, butyl rubber, nitrile rubber, neoprene rubber, natural rubber, polyethylene, PVC or viton; shoes plus socks.

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product through any type of irrigation system.

**FOR USE ON NONBEARING CITRUS (in greenhouse, shadehouse, field grown nurseries, solid settings, and as non-bearing resets)**

**NON-BEARING CITRUS**

**CITRUS ROOT WEEVIL COMPLEX:** For control of citrus root weevil complex including *Diaprepes abbreviatus*, the West Indian sugarcane rootstalk borer weevil, apply 20 ounces of MICROMITE 25WS per acre. Applications may be made by ground or aerial application equipment. Apply MICROMITE 25WS to newly expanded flush on citrus and/or when adult weevils are present. Use sufficient spray volume for adequate coverage of leaf surfaces. Repeat applications as needed for subsequent leaf flushes. The addition of a spray oil, such as FC435-66, enhances coverage and penetration of MICROMITE 25WS into the adult weevils themselves. Also, oil will deter attachment of weevil egg masses to leaf surfaces. A maximum of six (6) applications may be made per year.

NON-BEARING RESETS - except where planted in solid blocks, applications to individual trees must be by directed spray application, using hand-gun or shielded applicators. Do not spray bearing trees or allow drift to bearing trees.

MICROMITE 25WS should be used in conjunction with IPM practices including the early detection of target insect populations, threshold treatment levels, cultural control practices, and other procedures to manage target pest populations. Refer to local extension or university personnel regarding recommended IPM practices.

MICROMITE 25WS will not kill adult weevils. The activity of MICROMITE 25WS is through ingestion or contact and will result in reduction of the reproductive potential of weevils. It prevents eggs from hatching, thus preventing larvae from entering soil and feeding on citrus tree roots.

**LEPIDOPTEROUS LEAFMINERS:** For control of citrus lepidopterous leafminers (e.g. citrus leafminer) apply 20 ounces of MICROMITE 25WS per acre when oviposition begins on new growth flush. Applications may be made by ground or aerial application equipment. Use sufficient spray volume for adequate coverage of leaf surfaces. Repeat applications as needed for subsequent leaf flushes. Addition of FC435-66 enhances coverage and may enhance control of citrus leafminers. A maximum of six (6) applications may be made per year.

NON-BEARING RESETS - except where planted in solid blocks, applications to individual trees must be by directed spray application, using hand-gun or shielded applicators. Do not spray bearing trees or allow drift to bearing trees.

MICROMITE 25WS will not kill adult stages of leafminers. MICROMITE 25WS has activity on eggs, larval and pupal stages.

**WORKER PROTECTION:** Do not enter or allow worker entry into treated areas during the restricted entry interval (REI) of 12 hours. For all other worker protection standard requirements, such as personal protective clothing, refer to Standard Federal label for this product.

**Follow all applicable directions, restrictions, and precautions on the EPA registered label.  
This label must be in the possession of the user at the time of the pesticide application.**

001

MICROMITE® is a Registered Trademark of Uniroyal Chemical Company Inc., Middlebury, CT 06749  
DISTRIBUTED BY 24c (SLN) REGISTRANT:  
UNIROYAL CHEMICAL COMPANY, Inc.  
Middlebury, CT 06749

EPA SLN. NO. FL-010010