



Section 24(c) Special Local Need Label

FOR DISTRIBUTION AND USE ONLY WITHIN THE STATE OF WASHINGTON

FUSILADE® DX Herbicide

**EPA REG. NO. 100-1070
EPA SLN NO. WA-040007**

For Control of Annual and Perennial Grassy Weeds Including Quackgrass In Alfalfa (Grown For Seed)

FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR PEST CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.

DIRECTIONS FOR USE

It is a violation of federal law to use this product in a manner inconsistent with its labeling.

Follow all applicable directions, restrictions, and precautions including statements pertaining to the Worker Protection Standards, on the EPA-registered Fusilade DX Herbicide label.

This label must be in the possession of the user at the time of application.

Applications can be made by ground or air using the rates and timings given below:

WEED	APPLICATION	HEIGHT (inches)	RATE (fl. oz./A)
Bermudagrass ¹	1st	4-8	12
	2nd	4-8	8
Quackgrass ²	1st	6-10	12
	2nd	up to 10	8
Rhizome johnsongrass ³	1st	8-18	12
	2nd	6-12	8
Wirestem muhly ⁴	1st/2nd	4-12	12

¹Make second application to Bermudagrass if regrowth occurs (usually about 4 weeks after first application). To improve coverage, make applications in a minimum of 15 gallons per acre.

²Make second application 2-3 weeks after the first, but before the quackgrass exceeds 10 inches in height. Always use 1% v/v crop oil concentrate. Spot treatment is not recommended.

³Make first application before the boot stage.

⁴Make second application if regrowth occurs.

Environmental Hazards

This pesticide is toxic to fish. Fusilade DX should not be used under this SLN label where impact on listed threatened or endangered species is likely. You may refer to the WSDA Endangered Species Program website at <http://agr.wa.gov/PestFert/EnvResources/EndangSpecies.htm>, or contact the Washington Department of Fish & Wildlife, National Marine Fisheries Service (NOAA Fisheries) or US Fish & Wildlife Service for information regarding aquatic species listed as threatened or endangered. Consult the federal label for additional restrictions and precautions to protect aquatic organisms.

RESTRICTIONS:

1. All alfalfa seed screenings shall be disposed of in such a way that they cannot be distributed or used for food or feed. The seed conditioner shall keep records of screening disposal for three years from the date of disposal and shall furnish the records to the WSDA forthwith upon request. Disposal records shall consist of documentation from a controlled dumpsite, incinerator, or other equivalent disposal site and shall show the lot numbers, amount of material disposed of, its grower(s), and the date of disposal.
2. No portion of the alfalfa seed plant, including but not limited to green chop, hay, pellets, meal, whole seed, cracked seed, roots, bulbs, leaves and seed screenings may be used or distributed for food or feed purposes.
3. Alfalfa seed shall bear a tag or container label which forbids use of the seed for human consumption or animal feed.
4. Alfalfa seed may not be distributed for human consumption or animal feed.

USE PRECAUTIONS/RESTRICTIONS:

- Always add a crop oil concentrate at 0.5-1.0% or nonionic surfactant at 0.25-0.50% of the finished spray volume unless specified otherwise.
- Make aerial or ground applications when wind velocity favors on target product deposition (approximately 3-10 mph). Do not apply when wind velocity exceeds 10 mph.
- Do not plant rotational grass crops such as corn, sorghum and cereals within 60 days after last application.

WSDA Container Disposal Guidance: Pesticide containers must be properly cleaned prior to disposal. The best time to clean empty pesticide containers is during mixing and loading, because residue can be difficult to remove after it dries. Triple rinse (or pressure rinse) the pesticide container, empty all pesticide rinse water into the spray tank, and apply to a labeled crop or site. Recycling cleaned containers is the best method of container disposal. Information regarding the recycling of empty and cleaned plastic pesticide containers in Washington is available on the internet from WSU at <http://pep.wsu.edu/waste/wd.html> or from WSDA at <http://agr.wa.gov/PestFert/Pesticides/WastePesticide.htm>. Cleaned containers may also be disposed of in a sanitary landfill, if permitted by the county. Burning is not a legal method of container disposal in Washington.

Chemigation: For use under this SLN label do not apply this product through any type of irrigation system.

This label for Fusilade DX expires and must not be distributed or used in accordance with this SLN registration after December 31, 2009.

24(c) registrant:
Syngenta Crop Protection, Inc.
P.O. Box 18300
Greensboro, NC 27419-8300

EPA SLN No.: WA-040007

Label Code: WA1070065AA0504

Word: WAFusiladeDXAlfalfaforSeedSLNLb15-5-04