



FOR CONTROL OF CERTAIN WEEDS IN PEANUTS AND SOYBEANS

Active Ingredient	By Wt.
Flumioxazin*	51%
Other Ingredients	49%
Total	100%

* 2-[7-fluoro-3,4-dihydro-3-oxo-4-(2-propynyl)-2H-1,4-benzoxazin-6-yl]-4,5,6,7-tetrahydro-1H-isoindole-1,3(2H)-dione

Chateau™ Herbicide SW is a water dispersible granule containing 51% active ingredient.

EPA Reg. No. 59639-99

EPA Est. 11773-IA-01

KEEP OUT OF REACH OF CHILDREN

CAUTION

PRECAUTIONARY STATEMENTS

HAZARDS TO HUMANS & DOMESTIC ANIMALS

CAUTION

Harmful if inhaled or absorbed through the skin. Causes moderate eye irritation. Avoid breathing dust and spray mist. Avoid contact with skin, eyes or clothing.

FIRST AID	
If inhaled:	Move person to fresh air. If person is not breathing, call 911 or an ambulance, then give artificial respiration, preferably by mouth-to-mouth if possible. Call a poison control center or doctor for further treatment advice.
If on skin or clothing:	Take off contaminated clothing. Rinse skin immediately with plenty of water for 15-20 minutes. Call a poison control center or doctor for treatment advice.
If in eyes:	Hold eye open and rinse slowly and gently with water for 15-20 minutes. Remove contact lenses, if present, after the first 5 minutes, then continue rinsing eye. Call a poison control center or doctor for treatment advice.

(Continued)

FIRST AID (Continued)

If swallowed: Call a poison control center or doctor immediately for treatment advice.
Have person sip a glass of water if able to swallow.
Do not induce vomiting unless told to do so by the poison control center or doctor.
Do not give anything by mouth to an unconscious person.

HOT LINE NUMBER

Have the product container or label with you when calling a poison control center or doctor or going for treatment. You may also contact **1-800-892-0099** for emergency medical treatment information.

PERSONAL PROTECTIVE EQUIPMENT (PPE)

Some of the materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category A on an EPA chemical-resistance category selection chart.

Applicators and other handlers must wear: long-sleeved shirt and long pants, chemical-resistant gloves made of any waterproof material such as polyethylene or polyvinyl chloride, shoes and socks.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This product is toxic to non-target plants and aquatic invertebrates. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff may be hazardous to non-target plants and aquatic organisms in neighboring areas. Do not apply where runoff is likely to occur. Do not apply when weather conditions favor drift from treated areas. Do not contaminate water when disposing of equipment washwaters.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling.

READ ENTIRE LABEL. USE STRICTLY IN ACCORDANCE WITH PRECAUTIONARY STATEMENTS AND DIRECTIONS, AND WITH APPLICABLE STATE AND FEDERAL REGULATIONS.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific statements on this label about personal protective equipment (PPE), and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil or water, is: coveralls, chemical resistant gloves made of waterproof material, shoes plus socks.

DISCLAIMER, RISKS OF USING THIS PRODUCT, LIMITED WARRANTY AND LIMITATION OF LIABILITY

IMPORTANT: Read the entire Label including this Disclaimer, Risks of Using this Product, Limited Warranty and Limitation of Liability before using this product. If the terms are not acceptable THEN DO NOT USE THE PRODUCT; rather, return the unopened product within 15 days of purchase for a refund of the purchase price.

RISKS OF USING THIS PRODUCT

The Buyer and User (referred to collectively herein as "Buyer") of this product should be aware that there are inherent unintended risks associated with the use of this product which are impossible to eliminate. These risks include, but are not limited to, injury to plants and crops to which this product is applied, lack of control of the target pests or weeds, resistance of the target pest or weeds to this product, injury caused by drift, and injury to rotational crops caused by carryover in the soil. Such risks of crop injury, non-performance, resistance or other unintended consequences are unavoidable and may result because of such factors as weather, soil conditions, disease, moisture conditions, irrigation practices, condition of the crop at the time of application, presence of other materials either applied in the tank mix with this product or prior to application of this product, cultural practices or the manner of use or application (or a combination of such factors) all of which are factors beyond the control of Valent. The Buyer should be aware that these inherent unintended risks may reduce the harvested yield of the crop in all or a portion of the treated acreage, or otherwise affect the crop such that additional care, treatment and expense are required to take the crop to harvest. If the Buyer chooses not to accept these risks, THEN THIS PRODUCT SHOULD NOT BE APPLIED. By applying this product Buyer acknowledges and accepts these inherent unintended risks AND AGREES THAT ALL SUCH RISKS ASSOCIATED WITH THE APPLICATION AND USE ARE ASSUMED BY THE BUYER.

Valent shall not be responsible for losses or damages (including, but not limited to, loss of yield, increased expenses of farming the crop or such incidental, consequential or special damages that may be claimed) resulting from use of this product in any manner not set forth on the label. Buyer assumes all risks associated with the use of this product in any manner or under conditions not specifically directed or approved on the label.

(Continued)

LIMITED WARRANTY

Valent warrants only that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the label, under average use conditions, when used strictly in accordance with the label and subject to the Risks of Using This Product as described above. EXCEPT AS SET FORTH ABOVE, VALENT MAKES NO OTHER WARRANTIES, EITHER EXPRESSED OR IMPLIED. No agent or representative of Valent or Seller is authorized to make or create any other express or implied warranty.

LIMITATION OF LIABILITY

In no event shall Valent or Seller be liable for any incidental, consequential, indirect or special damages resulting from the use or handling of this product. The limitation includes, but is not limited to, loss of yield on all or any portion of the treated acreage, increased care, treatment or other expenses required to take the crop to harvest, increased finance charges or altered finance ratings, emotional or mental distress and/or exemplary damages.

THE EXCLUSIVE REMEDY OF THE BUYER, AND THE EXCLUSIVE MAXIMUM LIABILITY OF VALENT OR SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT SHALL BE THE RETURN OF THE PURCHASE PRICE OF THIS PRODUCT OR, AT THE ELECTION OF VALENT OR SELLER, THE REPLACEMENT OF THIS PRODUCT.

PROMPT NOTICE OF CLAIM

Valent must be provided notice as soon as Buyer has reason to believe it may have a claim, but in no event later than twenty-one days from date of planting, or twenty-one days from the date of application, whichever is latter, so that an immediate inspection of the affected property and growing crops can be made.

If Buyer does not notify Valent of any claims, in such period, it shall be barred from obtaining any remedy.

NO AMENDMENTS

Valent and Seller offer this product, and Buyer accepts it, subject to the foregoing Disclaimer, Risks of Using This Product, Limited Warranty and Limitation of Liability, which may not be modified by any oral or written agreement.

TANK MIXES

NOTICE: Tank mixing or use of this product with any other product which is not specifically and expressly authorized by the label shall be the exclusive risk of user, applicator and/or application advisor.

Read and follow the entire label of each product to be used in the tank mix with this product.

TABLE OF CONTENTS

GENERAL INFORMATION

- General Restrictions and Limitations
- Environmental Conditions and Biological Performance
 - Preemergence Applications
 - Burndown Applications
- Soil Characteristics
- Herbicide Rate
 - Residual Weed Control (Including Preemergence Applications or Applications As Part of a Spring Burndown Program)

Carrier Volume and Spray Pressure
 Preemergence Applications
 Burndown Applications
 Additives
 Burndown Applications
 Jar Test to Determine Compatibility of Adjuvants and
Chateau SW
 Sprayer Preparation
 Mixing Instructions
 Sprayer Cleanup
 Application Equipment
 Broadcast Application
 Band Application
 Aerial Application
 Crop Failure
 Rotational Restrictions
 Resistance Management

**DIRECTIONS FOR USE IN FALL AND SPRING
 PREPLANT BURNDOWN PROGRAMS IN PEANUTS
 AND SOYBEANS (Preemergence to Crop)**

General Restrictions and Limitations
 Fall Burndown Programs
 Weeds Controlled by Fall and Spring Preplant
 Burndown Programs.....Table 1
 Spring Burndown Programs

**DIRECTIONS FOR USE IN FALL AND SPRING
 BURNDOWN PROGRAMS IN COTTON, FIELD CORN,
 RICE, SORGHUM, SUGARCANE, SUNFLOWERS,
 TOBACCO AND WHEAT (Preemergence to Crop)**

General Restrictions and Limitations
 Fall Burndown Programs
 Spring Burndown Programs

**RECOMMENDATIONS FOR WEED CONTROL IN
 PEANUTS AND SOYBEANS**

Broadleaf Weeds Controlled by Residual Activity
 of *Chateau SW* in Peanuts and SoybeansTable 2
 Weeds Suppressed by Residual Activity of
Chateau SW in Peanuts and SoybeansTable 3

DIRECTIONS FOR USE IN PEANUTS

General Restrictions and Limitations
 Wind Management
 Timing to Peanuts
 Timing to Weeds
 Burndown - Preemergence to Peanuts,
 Postemergence to Weeds
 Additional Residual Grass Control: Sequential
 Additional Residual Grass Control: Tank Mixed

DIRECTIONS FOR USE IN SOYBEANS

General Restrictions and Limitations
 Timing to Soybeans
 Timing to Weeds
 Burndown - Preemergence to Soybeans,
 Postemergence to Weeds
 Increasing Speed of Glyphosate Burndown Activity
 Tank Mixes
 Tank Mix Partners for Control of Emerged Weeds
 in Reduced Tillage SoybeansTable 4
 Additional Residual Broadleaf Control
 Additional Residual Grass Control
 Roundup Ready® Program

GENERAL INFORMATION

Do not apply this product through any type of irrigation system.

AVOIDING SPRAY DRIFT AT THE APPLICATION SITE IS THE RESPONSIBILITY OF THE APPLICATOR.

The interaction of many equipment and weather related factors determine the potential for spray drift. The applicator is responsible for considering all these factors when making decisions. Where states have more stringent regulations, they should be observed.

Chateau SW provides residual control of susceptible weeds in peanuts and soybeans. *Chateau SW* can also be used as part of a burndown program in peanuts and soybeans. *Chateau SW* can be used as part of early preplant burndown programs in cotton, field corn, rice, sorghum, sugarcane, sunflowers, tobacco and wheat when applied more than 30 days prior to planting.

GENERAL RESTRICTIONS AND LIMITATIONS

- Do not apply this product when weather conditions favor spray drift from treated areas.
- Do not graze treated fields or feed treated forage or hay to livestock.
- Do not incorporate into the soil after application.
- When applying by air, observe drift management restrictions and precautions listed under "AERIAL APPLICATION".

ENVIRONMENTAL CONDITIONS AND BIOLOGICAL PERFORMANCE

PREEMERGENCE APPLICATIONS (Conventional Tillage)

Important: Crop injury may occur from applications made to poorly drained soils and/or applications made under cool, wet conditions. Risk of crop injury can be minimized by using on well drained soils, planting at least 1.5 inches deep, using high quality seed and completely covering seeds with soil prior to preemergence applications. Treated soil that is splashed onto newly emerged crops may result in temporary crop injury.

Moisture is necessary to activate *Chateau SW* in soil for residual weed control. Dry weather following applications of *Chateau SW* may reduce effectiveness. However, when adequate moisture is received after dry conditions, *Chateau SW* will control susceptible germinating weeds. *Chateau SW* may not control weeds that germinate after application but before an activating rainfall/irrigation or weeds that germinate through cracks resulting from dry soil.

When adequate moisture is not received after a *Chateau SW* application, weed control may be improved by irrigation with at least 1/4 inch of water. If emerged weeds are controlled by cultivation, residual weed control will be reduced.

BURNDOWN APPLICATIONS

For best results, *Chateau SW* should be applied as part of a burndown program to actively growing weeds. Applying *Chateau SW* under conditions that do not promote active weed growth will reduce herbicide effectiveness. Do not apply *Chateau SW* when weeds are under stress due to drought, excessive water, extremes in temperature, disease or low humidity. Weeds under stress tend to become less susceptible to herbicidal action. *Chateau SW* is most effective when applied under warm sunny conditions.

Reduced residual weed control may occur when burndown applications are made to fields where heavy crop and/or weed residue exist.

Chateau SW is rainfast one hour after application. Applications should not be made if rain is expected within one hour of application or efficacy may be reduced.

SOIL CHARACTERISTICS

Application of *Chateau* SW to soils with high organic matter and/or high clay content may require higher dosages than soils with low organic matter and/or low clay content. Application to cloddy seedbeds can result in reduced weed control.

HERBICIDE RATE

RESIDUAL WEED CONTROL (INCLUDING PREEMERGENCE APPLICATIONS OR APPLICATIONS AS PART OF A SPRING BURNDOWN PROGRAM)

Based upon soil characteristics (organic matter content and texture), the most difficult-to-control weed species being targeted, and the crop being grown, select the proper *Chateau* SW dosage from Table 2 when applying in peanuts or soybeans. Table 3 lists weeds that are suppressed by *Chateau* SW rates used in peanuts and soybeans.

CARRIER VOLUME AND SPRAY PRESSURE (Ground Equipment Only. See information for aerial equipment under "AERIAL APPLICATION".

PREEMERGENCE APPLICATIONS (Conventional Tillage)

To ensure uniform coverage, use 10 to 30 gals. spray solution per acre for conventional-tillage applications. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for preemergence herbicide application.

BURNDOWN APPLICATIONS (Prior to Crop Emergence)

To ensure thorough coverage in burndown applications, use 15 to 30 gals. spray solution per acre. Use 20 to 30 gals. per acre if dense vegetation or heavy crop residue is present. Nozzle selection should meet manufacturer's gallonage and pressure recommendations for postemergence herbicide application.

ADDITIVES

BURNDOWN APPLICATIONS (Prior to Crop Emergence)

Postemergence control of weeds from *Chateau* SW requires the addition of an agronomically approved adjuvant to the spray mixture. Either a crop oil concentrate or methylated seed oil which contains at least 15% emulsifiers and 80% oil may be used when applying *Chateau* SW as part of a burndown program. Some tank mix partners, such as Roundup UltraMax™, are formulated with sufficient adjuvants and do not require the addition of a crop oil concentrate or methylated seed oil when tank mixed with *Chateau* SW. The addition of a crop oil concentrate or methylated seed oil may increase the burndown activity on certain weeds such as cutleaf eveningprimrose and Carolina geranium. Mixing compatibility qualities should be verified by a jar test.

A spray grade nitrogen source (either ammonium sulfate at 2.0 to 2.5 lbs./A or a 28 to 32% nitrogen solution at 1 to 2 qts./A) may be added to the spray mixture along with either a crop oil concentrate or methylated seed oil to enhance weed control. The addition of a nitrogen source does not replace the need for crop oil concentrate or methylated seed oil.

JAR TEST TO DETERMINE COMPATIBILITY OF ADJUVANTS AND CHATEAU HERBICIDE SW

When using *Chateau* SW and an adjuvant, such as in stale seed bed or reduced tillage situations, a jar test should be performed before mixing commercial quantities of *Chateau* SW, when using *Chateau* SW for the first time, when using new adjuvants, or when a new water source is being used.

1. Add 1 pt. of the water to a quart jar. The water should be from the same source and temperature as which will be used in the spray tank mixing operation.
2. Add 1 gram of *Chateau* SW to the quart jar for every 3 oz. of *Chateau* SW/A being applied (1 gram if 3 oz./A is the desired *Chateau* SW rate), gently mix until product goes into suspension.
3. Add 60 ml (4 tbsp. or 2 fl. oz.) of the crop oil or methylated seed oil to the quart jar, gently mix.
4. If nitrogen is being used, add 16 ml (1 tbsp. or 0.5 oz.) of the 28 to 32% nitrogen source to the quart jar. If ammonium sulfate is being used, add 19 grams AMS to the quart jar in place of the 28 to 32% nitrogen.
5. Place cap on jar, invert 10 times, let stand for 15 minutes, evaluate.
6. An ideal tank mix combination will be uniform and free of suspended particles. If any of the following conditions are observed the choice of adjuvant should be questioned:
 - (1) Layer of oil or globules on the mixture's surface.
 - (2) Flocculation: fine particles in suspension or as a layer on the bottom of the jar.
 - (3) Clabbering: Thickening texture (coagulated) like gelatin.

SPRAYER PREPARATION

Before applying *Chateau* SW, start with clean, well maintained application equipment. The spray tank, as well as all hoses and booms, must be cleaned to ensure no residue from the previous spraying operation remains in the sprayer. Some pesticides, including but not limited to, the sulfonylurea and phenoxy herbicides (e.g., Classic® and 2,4-D, respectively), are active at very small amounts and can cause crop injury when applied to susceptible crops. The spray equipment must be cleaned according to the manufacturer's directions for the last product used before the equipment is used to apply *Chateau* SW. If two or more products were tank mixed prior to *Chateau* SW application, the most restrictive cleanup procedure should be followed.

MIXING INSTRUCTIONS

1. Fill clean spray tank 1/2 to 2/3 of desired level with clean water.
2. To ensure a uniform spray mixture, pre-slurry the required amount of *Chateau* SW with water prior to addition to the spray tank. Use a minimum of 1 gal. of water per 10 oz. of *Chateau* SW.
3. While agitating, slowly add the pre-slurried *Chateau* SW to the spray tank. Agitation should create a rippling or rolling action on the water surface.
4. If tank mixing *Chateau* SW with other labeled herbicides, add water soluble bags first, followed by dry formulations, flowables, emulsifiable concentrates, and then solutions. Prepare no more spray mixture than is required for the immediate spray operation.
5. Add any required adjuvants.
6. Fill spray tank to desired level with water. **Agitation should continue until spray solution has been applied.**
7. Mix only the amount of spray solution that can be applied the day of mixing. *Chateau* SW should be applied within 6 hours of mixing.

SPRAYER CLEANUP

Spray equipment must be cleaned each day following *Chateau SW* application. After *Chateau SW* is applied, the following steps must be used to clean the spray equipment.

1. Completely drain the spray tank, rinse the sprayer thoroughly, including the inside and outside of the tank and all in-line screens.
2. Fill the spray tank with clean water and flush all hoses, booms, screens and nozzles.
3. Top off tank, add 1 gal. of 3% household ammonia for every 100 gals. of water, circulate through sprayer for 5 minutes, and then flush all hoses, booms, screens and nozzles for a minimum of 15 minutes.
4. Drain tank completely.
5. Add enough clean water to the spray tank to allow all hoses, booms, screens and nozzles to be flushed for 2 minutes.
6. Remove all nozzles and screens and rinse them in clean water.

Spray equipment, including all tanks, hoses, booms, screens and nozzles, should be thoroughly cleaned before it is used to apply postemergence pesticides. Equipment with *Chateau SW* residue remaining in the system may result in crop injury to the subsequently treated crop.

APPLICATION EQUIPMENT

Application equipment should be clean and in good repair. Nozzles should be uniformly spaced on boom and frequently checked for accuracy.

BROADCAST APPLICATION

Apply *Chateau SW* and *Chateau SW* tank mixes with ground equipment using standard commercial sprayers equipped with flat fan or flood nozzles (preemergence applications only) designed to deliver the desired spray pressure and spray volume.

BAND APPLICATION

When banding, use proportionately less water and *Chateau SW* per acre.

AERIAL APPLICATION

Spray drift away from the site of application may cause damage to non-target vegetation. To minimize drift, apply the largest droplet size consistent with uniform coverage and satisfactory weed control. To obtain satisfactory application and avoid drift, the following directions must be observed:

- Do not apply during low-level inversion conditions, when winds are gusty or under other conditions that favor drift. Do not spray when wind velocity is less than 2 mph or more than 10 mph.
- Do not apply this product by air within 40 feet of non-target plants including non-target crops.
- Do not apply this product by air within 40 feet of streams, wetlands, marshes, ponds, lakes and reservoirs.
- **Carrier Volume and Spray Pressure:** When used as part of a burndown weed control program, apply *Chateau SW* in 7 to 10 gals. of water per acre. Application at less than 7 gals. per acre may provide inadequate control. When used for preemergence weed control, apply *Chateau SW* in 5 to 10 gals. per acre of water. The higher gallonage applications generally afford more consistent weed control. Do not exceed the nozzle manufacturer's recommended pressures. For many nozzle types, lower pressure produces larger droplets. When higher flow rates are needed, use higher flow rate nozzles instead of increasing pressure.

- **Nozzle Selection and Orientation:** Formation of very small drops may be minimized by appropriate nozzle selection, by orienting nozzles away from the air stream as much as possible and by avoiding excessive spray pressure. Use nozzles which produce flat or hollow cone spray patterns. Use non-drip type nozzles, such as diaphragm-type nozzles, to avoid unwanted discharge of spray solution. The nozzles must be directed toward the rear of the aircraft, at an angle between 0° and 15° downward. Do not place nozzles on the outer 25% of the wings or rotors.
- **Adjuvants and Drift Control Additives:** Refer to tank mix partner's label for adjuvant recommendation. Drift control additives may be used. When a drift control additive is used, read and carefully observe the cautionary statements and all other information appearing on the additive label.

CROP FAILURE

If the crop treated with *Chateau SW* is lost due to a catastrophe, such as hail or other forms of inclement weather, soybeans or peanuts can be replanted immediately, provided no more than 3 oz./A of *Chateau SW* has been used on the lost crop. Cotton, field corn, rice, sorghum, sugarcane, sunflowers, tobacco and wheat can be planted 30 days after a *Chateau SW* application, provided no more than 2 oz./A of *Chateau SW* had been used on the lost crop and a minimum of 1 inch of rainfall/irrigation has occurred between *Chateau SW* application and replanting. Crop injury may occur if these restrictions are not followed.

ROTATIONAL RESTRICTIONS

The following rotational crops may be planted after applying *Chateau SW* at the listed rate. Planting earlier than the recommended rotational interval may result in crop injury.

CHATEAU SW RATES	CROPS	ROTATION INTERVALS
2 oz./A or less	Peanut and Soybean	Immediately
	Cotton, Field Corn, Rice, Sorghum, Sunflower, Sugarcane, Tobacco and Wheat	30 days ¹
	Barley, Dry Bean, Field Pea, Rye and Sweet Corn	4 Months
Up to 3 oz./A	Alfalfa, Canola, Clover, Oats, Sugar Beet, and all other crops not listed	12 Months ²
	Peanut and Soybean	Immediately
	Cotton, Field Corn, Rice, Sorghum, Sunflower, Sugarcane, Tobacco and Wheat	2 Months
	Barley, Dry Bean, Field Pea, Rye and Sweet Corn	4 Months
	Alfalfa, Canola, Clover, Oats, Sugar Beet, and all other crops not listed	12 Months ⁽²⁾

1. At least one inch of rainfall/irrigation must occur between application and planting or crop injury may occur.
2. Successful soil bioassay must be performed prior to planting alfalfa, canola, sugar beets and other crops not listed.

RESISTANCE MANAGEMENT

Any weed population may contain or develop plants naturally resistant to herbicides in various mode of action classes. Resistant biotypes may eventually dominate the weed population if the same class of chemistry/mode of action herbicides are used repeatedly in the same field or in successive years. These resistant biotypes may not be adequately controlled by herbicides in a mode of action class for which resistance has developed. A gradual or total loss of weed control may occur over time. Other resistance mechanisms that are not linked to site of action, such as enhanced metabolism, may also exist. Appropriate resistance management strategies should be followed.

TO DELAY HERBICIDE RESISTANCE

- Avoid the use of herbicides that have a similar target site mode of action in consecutive years.
- Herbicide use should be based on an Integrated pest Management (IPM) program that includes scouting, record keeping, and consideration of cultivation practices, water management, weed free crop seed, crop rotation, and other chemical or cultural control practices.
- Monitor treated weed population for resistance development and report suspected resistance.
- Contact your local extension or crop expert (advisor) for any additional pesticide resistance management and/or IPM recommendations for specific crops and weed biotypes.
- For further information contact Valent U.S.A. Corporation at the following toll free number: 1-800-682-5368.

DIRECTIONS FOR USE IN FALL AND SPRING PREPLANT BURNDOWN PROGRAMS IN PEANUTS AND SOYBEANS (Preemergence to Crop)

GENERAL RESTRICTIONS AND LIMITATIONS

- Do not apply to frozen or snow covered soil.
- Do not perform any tillage operation after application or residual weed control will be reduced.

FALL BURNDOWN PROGRAMS

Chateau SW can be used in combination with labeled preplant burndown herbicides to control emerged weeds and provide residual weed control in fields that will be planted the following spring. Application must be made no earlier than October 15 in Region 2 or November 15 in Region 1 or when soil temperature falls below 50°F at a two inch depth to maintain residual weed control into the spring (April 1 in Region 1 and May 1 in Region 2) or up until planting, whichever comes first.

Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

Fall Application Regions:

Region 1: Alabama, Arkansas, Georgia, Louisiana, Kentucky, Mississippi, Oklahoma, Tennessee and Virginia.

Region 2: Delaware, Kansas, Illinois, Indiana, Iowa, Maryland, Michigan, Minnesota, Missouri, Nebraska, North Dakota, Ohio, Pennsylvania, South Dakota, West Virginia and Wisconsin.

Weeds controlled by postemergence or residual activity are listed in Table 1.

Preplant burndown treatment tank mixes and rates are:

Herbicide	Rate
Program 1	
<i>Chateau</i> SW plus	2 to 3 oz./A
Glyphosate plus	0.5 to 1.0 lb. ai/A (equivalent to 1 to 2 pt./A of Roundup Original™)
2,4-D LVE (2,4-D for use on preplant soybeans only) plus	0.5 to 1.0 lb. ai/A (equivalent to 1 to 2 pt./A of 2,4-D 4 LVE)
NIS + AMS or	0.5% v/v + 17 lbs./100 gals. of water
Program 2	
<i>Chateau</i> SW plus	2 to 3 oz./A
Glyphosate plus	0.5 to 1.0 lb. ai/A (equivalent to 1 to 2 pt./A of Roundup Original)
COC ¹ or	1 pt./A
NIS + AMS	0.5% v/v + 17 lbs./100 gals. of water
Program 3	
<i>Chateau</i> SW plus	2 to 3 oz./A
2,4-D LVE (2,4-D for use on preplant soybeans only) plus	0.5 to 1.0 lb. ai/A (equivalent to 1 to 2 pt./A of 2,4-D 4 LVE)
COC	1 pt./A

1. Crop oil concentrate has been found to increase glyphosate burndown on emerged cutleaf eveningprimrose or Carolina geranium.

Table 1. Weeds Controlled by Fall and Spring Preplant Burndown Programs

WEEDS CONTROLLED ¹		Postemergence			
COMMON NAME	SCIENTIFIC NAME	Program 1	Program 2	Program 3	Residual
		Weeds 3 Inches or Less			
Chickweed	<i>Stellaria media</i>	Yes	Yes	No	Yes
Common Mouseear	<i>Cerastium vulgatum</i>	Yes	Yes	No	Yes
Dandelion	<i>Taraxacum officinale</i>	Yes	No	Yes ²	Yes
Henbit	<i>Lamium amplexicaule</i>	Yes	Yes	Yes	Yes
Marestail/Horseweed	<i>Conya canadensis</i>	Yes	Yes	Yes	Yes
Groundsel, Cressleaf	<i>Senecio glabellus</i>	Yes	Yes	--	Yes
Purple Deadnettle	<i>Lamium purpureum</i>	Yes	Yes	Yes	Yes
		Weeds 12 Inches or Less			
Carolina Geranium	<i>Geranium carolinianum</i>	Yes	Yes	Yes	--
Eveningprimrose, Cutleaf ³	<i>Oenothera laciniata</i>	Yes	Yes	Yes	Yes
Mustard, Wild	<i>Brassica kabur</i>	Yes	Yes	Yes	Yes
Shepherds-purse	<i>Capsella bursa-pastoris</i>	Yes	Yes	Yes	Yes

1. Refer to glyphosate and/or 2,4-D labels for additional weeds controlled and rotational restrictions.
2. 1 lb. ai/A of 2,4-D LVE (equivalent to 2 pt./A of 2,4-D 4 LVE) should be used for control of emerged dandelion.
3. Program 1 should be used to control eveningprimrose that are nearing 12 inches in height or are past the rosette stage. Programs 2 or 3 should be used to control eveningprimrose that are 12 inches or less and in the rosette stage.

SPRING BURNDOWN PROGRAMS

Chateau SW can be used in combination with labeled preplant burndown herbicides to control emerged weeds and provide residual weed control prior to crop emergence. Weeds controlled by residual activity are listed in Table 2, Section A.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row. Apply *Chateau* SW after planting when these types of planters are used (within 3 days of planting and before the crop emerges).

Chateau SW can be used at 1 to 3 oz./A with labeled preplant burndown herbicides to enhance the speed of burndown and increase weed spectrum.

Chateau SW can be used at 1 to 3 oz./A in soybean and peanut burndown programs. See “DIRECTIONS FOR USE IN SOYBEANS”, “DIRECTIONS FOR USE IN PEANUTS” for more information.

DIRECTIONS FOR USE IN FALL AND SPRING BURNDOWN PROGRAMS IN COTTON, FIELD CORN, RICE, SORGHUM, SUGARCANE, SUNFLOWERS, TOBACCO AND WHEAT (Preemergence to Crop)

GENERAL RESTRICTIONS AND LIMITATIONS

- Do not apply to frozen or snow covered soil.
- Do not perform any tillage operation after application or residual weed control will be reduced.
- *Chateau* SW can be used at 1 to 2 oz./A with labeled burndown herbicides to enhance the speed of burndown and increase weed spectrum. A minimum of 30 days must pass, and 1 inch of rainfall/irrigation must occur, between *Chateau* SW application and planting of cotton, field corn, rice, sorghum, sugarcane, sunflowers, tobacco or wheat. Refer to most restrictive label for minimum interval between application and planting.

FALL BURNDOWN PROGRAMS

Chateau SW can be used in combination with labeled burndown herbicides to control emerged weeds and provide residual weed control in fields that will be planted the following spring. Application must be made no earlier than October 15 in Region 2 or November 15 in Region 1 or when soil temperature falls below 50°F at a two inch depth to maintain residual weed control into the spring (April 1 in Region 1 and May 1 in Region 2) or up until planting, whichever comes first.

Abnormally warm or wet winters will reduce the length of weed control observed in the spring.

SPRING BURNDOWN PROGRAMS

Chateau SW can be used in combination with labeled burndown herbicides to control emerged weeds and provide residual weed control prior to crop emergence. Weeds controlled by residual activity are listed in Table 2, Section A. Crops that will be planted following application must be in compliance with the rotational interval listed in the “Rotational Restrictions” table preceding.

No-till planters that incorporate the soil during planting may result in decreased weed control in the row.

RECOMMENDATIONS FOR WEED CONTROL IN PEANUTS AND SOYBEANS

Table 2 lists broadleaf weeds controlled by residual activity of *Chateau* SW in peanuts and soybeans. Table 3 lists weeds suppressed by residual activity of *Chateau* SW in peanuts and soybeans.

Table 2. Broadleaf Weeds Controlled by Residual Activity of *Chateau* Herbicide SW in Peanuts and Soybeans

BROADLEAF WEED SPECIES				
SECTION A				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	CHATEAU SW RATES
Carpetweed	<i>Mollugo verticillata</i>	Up to 5%	All Soil Types	2.0 oz./A
Chickweeds				
Common	<i>Stellaria media</i>			
Mouseear	<i>Cerastium vulgatum</i>			
Dandelion	<i>Taraxacum officinale</i>			
Eclipta	<i>Eclipta prostrata</i>			
Eveningprimrose, Cutleaf	<i>Oenothera laciniata</i>			
Florida Pusley	<i>Richardia scabra</i>			
Henbit	<i>Lamium amplexicaule</i>			
Kochia	<i>Kochia scoparia</i>			
Lambsquarters, Common	<i>Chenopodium album</i>			
Little Mallow	<i>Malva parviflora</i>			
Marestail/Horseweed	<i>Conyza canadensis</i>			
Nightshades				
Black	<i>Solanum nigrum</i>			
Eastern Black	<i>Solanum ptycanthum</i>			
Pigweeds				
Redroot	<i>Amaranthus retroflexus</i>			
Smooth	<i>Amaranthus hybridus</i>			
Spiny Amaranth	<i>Amaranthus spinosus</i>			
Tumble	<i>Amaranthus albus</i>			
Prickly Sida (Teaweed)	<i>Sida spinosa</i>			
Puncturevine	<i>Tribulus terrestris</i>			
Purslane, Common	<i>Portulaca oleracea</i>			
Redmaids	<i>Calandrinia ciliata</i> , var. <i>menziesii</i>			
Shepherds-purse	<i>Capsella bursa-pastoris</i>			
Smallflower Morningglory	<i>Jacquemontia tamnifolia</i>			
Spotted Spurge	<i>Euphorbia maculata</i>			
Venice Mallow	<i>Hibiscus trionum</i>			
SECTION B				
All weeds listed in Section A plus:				
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	SOIL TYPE	CHATEAU SW RATES
Coffee Senna	<i>Cassia occidentalis</i>	Up to 3%	All Soil Types	3.0 oz./A Peanuts ²
Common Ragweed ¹	<i>Ambrosia artemisiifolia</i>			
Golden Crownbeard	<i>Verbesina encelioides</i>			2.5 oz./A Soybeans
Florida Beggarweed	<i>Desmodium tortuosum</i>			
Hairy Indigo	<i>Indigofera hirsuta</i>			
Hemp Sesbania	<i>Sesbania exaltata</i>	3 to 5%	Coarse and Medium Soils: (sandy loam, loamy sand, loamy, silt-loam, silt, sandy clay, sandy clay loam)	3.0 oz./A Peanuts ²
Jimsonweed	<i>Datura stramonium</i>			
Morningglories ³				2.5 oz./A Soybeans
Entireleaf	<i>Ipomoea hederacea</i> var. <i>integruscula</i>			
Ivyleaf		3 to 5%	Fine Soils: (silty clay, silty clay loam, clay, clay loam)	3.0 oz./A Peanuts ² and Soybeans
Red/Scarlet	<i>Ipomoea hederacea</i>			
Tall	<i>Ipomoea coccinea</i>			
Mustard, Wild	<i>Ipomoea purpurea</i>			
Palmer Amaranth	<i>Brassica kaber</i>			
Spurred Anoda	<i>Amaranthus palmeri</i>			
Tropic Croton	<i>Anoda cristata</i>			
Waterhemp ¹	<i>Croton glandulosus</i>			
Common				
Tall	<i>Amaranthus rudis</i>			
Wild Poinsettia	<i>Amaranthus tuberculatus</i>			
	<i>Euphorbia heterophylla</i>			

1. A postemergence herbicide, such as Cobra[®], Phoenix[™] or glyphosate (Roundup Ready soybeans only) may be needed following a preemergence application of *Chateau* SW to adequately control common ragweed or waterhemp in soybean fields with heavy pressure.
2. Due to differences in crop canopy timing between peanuts and soybeans, 3.0 oz./A of *Chateau* SW should be used in peanuts, regardless of soil type and organic matter content, except in the states of Virginia, North Carolina and Oklahoma, where a maximum of 2 oz./A can be applied in peanuts.
3. Morningglory species are not adequately controlled on fine soils or soils with greater than 3% organic matter.

Table 3. Weeds Suppressed by Residual Activity of *Chateau* Herbicide SW in Peanuts and Soybeans

BROADLEAF WEED SPECIES			
COMMON NAME	SCIENTIFIC NAME	ORGANIC MATTER	OUNCES PER ACRE
Bristly Starbur	<i>Acanthospermum hispidum</i>	Up to 5%	2.5 to 3.0
Copperleaf, Hophornbeam	<i>Acalypha ostryifolia</i>		
Ragweed, Giant	<i>Ambrosia trifida</i>		
Russian Thistle	<i>Salsola iberica</i>		
Smartweeds			
Ladysthumb	<i>Polygonum persicaria</i>		
Pennsylvania	<i>Polygonum pennsylvanicum</i>		
Velvetleaf	<i>Abutilon theophrasti</i>		
GRASS WEED SPECIES			
Barnyardgrass	<i>Echinochloa crus-galli</i>		
Crabgrass, Large	<i>Digitaria sanguinalis</i>		
Foxtail, Giant	<i>Setaria faberi</i>		
Goosegrass	<i>Eleusine indica</i>		
Lovegrass, California	<i>Eragrostis diffusa</i>		
Panicums			
Fall	<i>Panicum dichotomiflorum</i>		
Texas	<i>Panicum texanum</i>		
Signalgrass, Broadleaf	<i>Brachiaria platyphylla</i>		

DIRECTIONS FOR USE IN PEANUTS

GENERAL RESTRICTIONS AND LIMITATIONS

- Do not apply more than 3 oz. of *Chateau* SW per acre during a single growing season.
- Do not apply more than 2 oz./A in the states of Virginia, North Carolina or Oklahoma, where climatic conditions may result in unacceptable injury to peanuts.
- Do not use on peanuts grown for seed.
- Do not irrigate when peanuts are cracking.
- Do not tank mix with Strongarm®.
- Do not use on the following peanut varieties: Perry or NC 10C.

WIND MANAGEMENT

In areas where shallow cultivation is used between rows to reduce wind-borne sand damage to peanuts, weed control from *Chateau* SW may be reduced.

TIMING TO PEANUTS

Chateau SW may be applied to peanuts prior to planting or preemergence (after planting). Preemergence applications of *Chateau* SW must be made within 2 days after planting and prior to peanut emergence. Application after the peanuts have begun to crack, or are emerged, will result in severe crop injury. Application should not be made when peanuts have begun to crack. Select *Chateau* SW rate from Table 2 according to anticipated weed spectrum.

TIMING TO WEEDS

BURNDOWN - PREEMERGENCE TO PEANUTS, POSTEMERGENCE TO WEEDS

Chateau SW, applied as part of a burndown program, may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where peanuts will be planted directly into a stale seedbed, cover crop, or in previous crop residues. Apply *Chateau* SW before planting, during planting, or after planting, but before the crop emerges. For control of emerged weeds, tank mix *Chateau* SW with glyphosate. Refer to glyphosate label for recommended rate and application pressure. To ensure thorough coverage, use a

minimum of 15 gals. of spray solution per acre. *Chateau* SW tank mixes applied to assist in the control of emerged weeds must be applied with an adjuvant, such as crop oil concentrate or methylated seed oil at 1 to 2 pt./A. A spray grade nitrogen source (either ammonium sulfate at 2.0 to 2.5 lbs./A or 28 to 32% nitrogen solution at 1 to 2 qts./A) may be added to increase herbicidal activity.

Preemergence (conventional tillage) applications of *Chateau* SW must be applied prior to weed emergence.

ADDITIONAL RESIDUAL GRASS CONTROL: SEQUENTIAL

Chateau SW may be applied sequentially following a preplant incorporated application of trifluralin (states of Texas, Oklahoma and New Mexico only), Sonalan®, Dual® (metolachlor), pendimethalin or Frontier®.

ADDITIONAL RESIDUAL GRASS CONTROL: TANK MIXED

Chateau SW can be tank mixed with alachlor, metolachlor, or Frontier for additional grass and broadleaf weed control. *Chateau* SW can also be tank mixed with pendimethalin or Sonalan in states where they are labeled, provided overhead irrigation guidelines on the pendimethalin and/or Sonalan labels are followed.

Read tank mix product label for rates and weeds controlled. Always read and follow label directions for all tank mix products before using. The most restrictive labeling of any tank mix product must be followed. *Chateau* SW, when applied according to label use directions, will control the weeds listed in Table 2. This label makes no claims concerning control of other weed species.

DIRECTIONS FOR USE IN SOYBEANS

GENERAL RESTRICTIONS AND LIMITATIONS

- Do not apply more than 3 oz. of *Chateau* SW per acre during a single growing season.
- Do not use *Chateau* SW in soybeans in the same field flufenacet (Axiom®, Domain®), alachlor (Micro-Tech®), metolachlor (Dual products or Boundary®), or dimethenamid (Frontier or Outlook®) will be used or soybean injury may occur.
- Do not irrigate when soybeans are cracking.

TIMING TO SOYBEANS

Chateau SW may be applied to soybeans prior to planting or preemergence (after planting). Preemergence application of *Chateau SW* must be made within 3 days after planting and prior to soybean emergence. Application after the soybeans have begun to crack, or are emerged, will result in severe crop injury. Application should not be made when soybeans have begun to crack. Select *Chateau SW* rate from Table 2 according to anticipated weed spectrum.

TIMING TO WEEDS

BURNDOWN - PREEMERGENCE TO SOYBEANS, POSTEMERGENCE TO WEEDS

Chateau SW, applied as part of a burndown program, may be used for residual weed control, as well as to assist in postemergence burndown of many annual and perennial weeds where soybeans will be planted directly into a stale seedbed, cover crop, or in previous crop residues. For control of emerged weeds, choose the most appropriate tank mix partner from Table 4. Apply *Chateau SW* with ground equipment before planting, during planting, or after planting, **but before the crop emerges**. To ensure thorough coverage, use a minimum of 15 gals. of spray solution per acre. Refer to tank mix partner's label for recommended application pressure. All *Chateau SW* tank mixes applied to assist in the control of emerged weeds must be applied with crop oil concentrate or methylated seed oil at 1 to 2 pt./A.

INCREASING SPEED OF GLYPHOSATE BURNDOWN ACTIVITY

Chateau SW, at rates as low as 1.0 oz./A, may be tank mixed with glyphosate (Roundup®) to increase the speed of burndown activity compared to glyphosate applied alone. Residual weed control will not be provided at rates lower than 2 oz./A; however, suppression of the weeds in Table 2, Section A may occur at *Chateau SW* rates as low as 1 oz./A.

TANK MIXES

Chateau SW may be tank mixed with the herbicides listed in Table 4 for increased burndown activity, additional residual broadleaf, and/or additional grass control. Refer to tank mix partner's label for adjuvant recommendations.

Table 4. Tank Mix Partners for Control of Emerged Weeds in Reduced Tillage Soybeans

TANK MIX PARTNERS	TARGET WEEDS ⁽¹⁾
Gramoxone® Extra	Annual Grasses Henbit
Glyphosate	General Burndown
Select® 2 EC	Annual Grasses
Scepter® 70 DG	Cocklebur Common Sunflower
2,4-D LVE	Marestail Giant Ragweed Dandelion

1. Refer to tank mix product labels for specific recommendations for control of emerged weeds present.

ADDITIONAL RESIDUAL BROADLEAF CONTROL

Chateau SW can be tank mixed with metribuzin, FirstRate®, Lorox®, Pursuit Plus®, Python®, Squadron®, Scepter or Steel® for additional broadleaf control.

ADDITIONAL RESIDUAL GRASS CONTROL

Chateau SW can be tank mixed with pendimethalin or Command® for additional grass control. Tank mixes with fluthiamide (Axiom or Domain), metolachlor (Dual products or Boundary), dimethenamid (Frontier or Outlook) or alachlor (Micro-Tech), may result in severe injury to soybeans when application is followed by prolonged periods of cool wet weather and should not be used with *Chateau SW*.

ROUNDUP READY PROGRAM

Chateau SW may be applied as part of a burndown program or preemergence in conventional tillage programs, at 2 to 3 oz./A to reduce early season weed competition from waterhemp, velvetleaf, nightshade and morningglories as well as other weeds listed in Tables 2 and 3 in Roundup Ready programs. A sequential post emergence application of glyphosate will be required to control weeds not controlled by *Chateau SW*.

Read tank mix product label for rates and weeds controlled. Always read and follow label directions for all tank mix products before using. The most restrictive labeling of any tank mix product must be followed. *Chateau SW*, when applied according to label use directions, will control the weeds listed in Table 2. This label makes no claims concerning control of other weed species.

STORAGE AND DISPOSAL

PROHIBITIONS

Do not contaminate water, food or feed by storage, disposal or cleaning of equipment.

STORAGE

Keep pesticide in original container.

Store in a cool, dry, secure place.

Do not put formulation or dilute spray solution into food or drink containers.

Do not contaminate food or foodstuffs.

Do not store or transport near feed or food.

Not for use or storage in or around the home.

For help with any spill, leak, fire or exposure involving this material, call day or night (800) 892-0099.

PESTICIDE DISPOSAL

Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL

Triple rinse (or equivalent). Do not reuse container. Offer for recycling or reconditioning, or puncture and dispose of in a sanitary landfill or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

Copyright © 2004 Valent U.S.A. Corporation

Axiom, Domain – Reg. TMs of Bayer

Boundary, Dual – Reg. TMs of Syngenta

Chateau, *Cobra*, *Phoenix*, *Products That Work*, *From People Who Care*, *Select* – TMs and Reg. TMs of Valent U.S.A. Corporation

Classic, Lorox – Reg. TMs of DuPont

Command – Reg. TM of FMC Corporation

FirstRate, Python, Sonalan, Strongarm – Reg. TMs of Dow AgroSciences LLC

Frontier, Outlook, Pursuit Plus, Scepter, Squadron, Steel – Reg. TMs of BASF Corporation

Gramoxone – Reg. TM of Zeneca Ag Products
Micro-Tech, Roundup, Roundup Original, Roundup Ready
and Roundup UltraMax – TMs and Reg. TMs of Monsanto
Company

Manufactured for:

Valent U.S.A. Corporation

P.O. Box 8025

Walnut Creek, CA 94596-8025

EPA Reg. No. 59639-99

EPA Est. 11773-IA-01

Form 1487-A

Made in U.S.A.

Information contained in this booklet is accurate at the time
of printing. Since product testing is a continuous process,
please read and follow the directions on the product label
for the most current directions and precautionary
statements.

Always check with your state to verify state registration
status.

Products That Work, From People Who Care™

2004-CHSW-0001 10/03 (AV)

Chateau Herbicide SW

For more information
Valent U.S.A. Corporation:
800-6-VALENT(682-5368)
www.valent.com

