

# CAUTION

**KEEP OUT OF REACH OF CHILDREN  
READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

## FLUPROMIX™ 500

## HERBICIDE

**ACTIVE CONSTITUENTS: 250 g/L FLUOMETURON  
250g/L PROMETRYN**

<b>GROUP</b>	<b>C</b>	<b>HERBICIDE</b>
--------------	----------	------------------

**For control of many broadleaf weeds and annual grasses in cotton.**

## 20, 200 LITRES

**SIPCAM PACIFIC AUSTRALIA PTY LTD  
Suite 11, 23-31 Gheringhap Street  
GEELONG VIC 3220  
ABN 94 073 176 888  
TEL: 03 5223 33746**

<sup>TM</sup> Trademark of Sipcam Pacific Australia Pty Ltd

**Directions for Use:****For use in cotton only or the control of those weeds listed in the General Instructions****RESTRAINTS**

DO NOT use on light, sandy soils or soils of low organic content.

<b>Method of Application</b>	<b>Rate per hectare</b>	<b>State</b>	<b>Critical Comments</b>
Soil incorporated, pre-sowing treatment followed (if required) by pre-emergence band treatment	<b>Pre-sowing</b> 2.5 to 5 litres  <b>Pre-emergence</b> 2 to 3 litres per sprayed hectare	NSW, Qld Only	Apply initial treatment just prior to incorporation. Lightly incorporate to a depth of 5 cm using suitable equipment to ensure accurate and even incorporation. Following sowing, apply a further pre-emergence band treatment of FLUPROMIX to compensate for the chemical scraped off the top of the hill by the planting operations. Soil should preferably be moist at the time of application and significant rainfall or irrigation is required within 3-5 days of application to incorporate the herbicide into the soil.
Pre-emergence treatment	3 to 5 litres	NSW, Qld only	Apply immediately after sowing, before weed or crop germination, as an overall or band treatment. The minimum band width should be 40 cm. Soil should preferably be moist at the time of application and significant rainfall or irrigation is required within 3-5 days of application to incorporate the herbicide into the soil.
Post-emergence early spraying	1.5 to 2.5 litres	Qld only	Apply as directed spray to avoid contact with the cotton foliage to control broadleaf weeds and grasses which may have escaped previous herbicide treatments. Apply when cotton is 30-50 cm high and weeds are not more than 8 cm high and actively growing. Add a non-ionic surfactant at the rate of 50 mL per 100 litres.
Post-emergence lay-by spraying	2 to 3.5 litres	NSW, Qld only	Apply as directed spray underneath cotton foliage in 100-200 litres of water per hectare after the last cultivation before new weed germinations occur, or to weeds not more than 8 cm high and actively growing. Add a non-ionic surfactant at the rate of 50 mL per 100 litres.

**NOT TO BE USED FOR ANY PURPOSE, OR ANY MANNER, CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION.****WITHHOLDING PERIOD: NOT REQUIRED WHEN USED AS DIRECTED**

## General Instructions

### Weeds Controlled

Flupromix 500 Herbicide controls weeds and grasses including Barley Grass, Barnyard Grass, Bathurst Burr, Bellvine, Black-berry Nightshade, Black Pigweed, Bladder Kemia, Boggabri Weed, Caltrop (Yellow Vine), Capeweed, Caustic Weed (Qld only), Common Sowthistle, Common Thornapple, Fat-hen, Liverseed Grass, Mossman River Grass, Noogoora Burr, Paterson's Curse, Peach Vine, Pigweed, Prickly Paddy Melon, Redroot Amaranth, Annual Ryegrass, Sesbania Pea, Slim Amaranth, Summer Grass, Variegated Thistle, Wild Gooseberry, Winter Grass, Yellowflower Devil's-claw.

### RESISTANT WEEDS WARNING

Flupromix 500 Herbicide is a member of the triazine and urea group of herbicides and has the inhibitor of photosynthesis at photosystem II mode of action. For weed resistance management this is a mode of action Group C herbicide.

Some naturally-occurring weed biotypes resistant to Flupromix 500 Herbicide and other inhibitor of photosynthesis at photosystem II herbicides may exist through normal genetic variability in any weed population. The resistant individuals can eventually dominate the weed population if these herbicides are used repeatedly. These resistant weeds will not be controlled by Flupromix 500 Herbicide or other inhibitor of photosynthesis at photosystem II herbicides.

Since the occurrence of resistant weeds is difficult to detect prior to use, Sipcarn Pacific Australia Pty Ltd accepts no liability for any losses that may result from the failure of Flupromix 500 Herbicide to control resistant weeds.

Advice as to strategies and alternative treatments that can be used should be obtained from your local supplier, consultant, local Department of Agriculture, Primary Industries Department or a Sipcarn representative.

### Mixing

Settling may occur after storage for some weeks. Stir product or invert container several times before opening. Pour the product into the spray vat through a strainer to remove any dry particles or flakes which can occasionally occur under hot storage conditions. Add the full quantity of Flupromix to the partly filled spray tank while agitating. Fill tank and agitate to ensure thorough mixing. Continue agitation while spraying. Agitate vigorously from the bottom if allowed to stand. Reseal part used container immediately. DO NOT allow partly filled spray tank to stand overnight.

### APPLICATION

**Boom Spray:** 100 to 200 litres per ha. **Aircraft:** minimum of 10 litres per ha. Add a non-ionic surfactant for post-emergence application only.

### Re-entry Period

DO NOT enter treated areas until spray has dried.

### Precaution

Severe injury to cotton plants may result if heavy rain occurs between pre-emergence application and crop emergence, or shortly after post-emergence spraying, and this risk must be assumed by the user.

DO NOT make more than one post-emergence application per season.

Crops other than cotton should not be planted within six months of applying Flupromix. Use the higher recommended rates on heavy clay soils or soils of high organic content and the lower recommended rates on medium soils.

Seed beds must be well prepared and as free as possible from trash and clods.

### **Protection of Crops, Native and other Non-target Plants**

DO NOT apply under weather conditions or from spraying equipment that may cause spray to drift over nearby susceptible plants or crops, cropping lands or pastures.

### **Protection of Wildlife, Fish, Crustaceans and Environment**

DO NOT contaminate streams, rivers or waterways with the product or used containers. Avoid use in channels and drains where roots or desirable plants may extend and in other situations where desirable trees grow close to treated areas.

### **Storage and Disposal**

Store in the closed, original container in a well-ventilated areas as cool as possible out of direct sunlight.

Triple or preferably pressure rinse containers before disposal. Add rinsings to spray tank. DO NOT dispose of undiluted chemicals on-site. If recycling, replace cap and return clean containers to recycler or designated collection point.

If not recycling, break, crush or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

### **SAFETY DIRECTIONS**

May irritate the eyes and skin. Avoid contact with eyes and skin. DO NOT inhale spray mist. If product gets on skin, immediately wash areas with soap and water. After use and before eating, drinking or smoking, wash hands, arms and face thoroughly with soap and water.

### **FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (phone 131 126).

### **Material Safety Data Sheet**

If additional hazard information is required, refer to the Material Safety Data Sheet.

### **NOTICE TO BUYER**

This product is not to be used for any purpose or in any way contrary to label instructions. The buyer by use or application of this product acknowledges so doing upon the terms and conditions agreed between Sipcam Pacific Australia Pty Limited and the buyer but not limited to the following. To the extent permitted by law all conditions and other warranties or other rights of action which buyer or any other user may have against Sipcam Pacific Australia Pty Limited (Sipcam) or Seller are hereby excluded. Sipcam hereby gives notice to buyer and other users that it will not accept responsibility for any indirect or consequential loss arising from reliance on product information or advice provided by Sipcam or on its behalf unless it can be established that such information or advice was provided negligently and that the product has been used strictly as directed. Notwithstanding the generality of this notice Sipcam will not be held liable for any loss, injury or damage indirect or consequential arising from the sale, supply, use or application of this product. Sipcam's liability shall in all circumstances be limited to replacement of the product or a refund of the purchase price paid therefor.

NRA Approval No: 53766/0601

Batch No:

Date of Manufacture: