

**POISON**  
**KEEP OUT OF REACH OF CHILDREN**  
**READ SAFETY DIRECTIONS BEFORE OPENING OR USING**

**Jockey<sup>®</sup>**

**SYSTEMIC SEED FUNGICIDE**

**Active Constituent: 167 g/L FLUQUINCONAZOLE**

**GROUP C FUNGICIDE**

**For the suppression of take-all and septoria tritici blotch, the early season control of leaf rust and stripe rust and control of common bunt, loose smut and flag smut in wheat. For the suppression of blackleg in canola.**

### GENERAL INSTRUCTIONS

Jockey is a novel quinazoline-based triazole fungicide for the protection of wheat against the root disease, take-all, and protection of canola against blackleg.

When Jockey is applied as a seed treatment to wheat, the development of take-all is suppressed. Suppression of take-all with Jockey is characterised by reduced root lesions and a reduced number of whiteheads.

Jockey also provides season long control of common bunt, seed- and soil-borne flag smut and loose smut in wheat. In addition Jockey provides early season control of leaf rust, stripe rust and suppression of septoria tritici blotch in wheat.

Suppression of septoria tritici blotch with Jockey results in a reduction in the visible symptoms of this disease.

When Jockey is applied as a seed treatment to canola, the development of blackleg disease is suppressed. Suppression of blackleg with Jockey is characterised by reduced leaf infection, reduced plant lodging and increased plant survival.

Jockey is applied to wheat and canola seed prior to sowing. The low water solubility of the active ingredient, fluquinconazole, allows extended uptake for long lasting protection of the plant, thus reducing the yield decline commonly associated with the above-mentioned root, grain and foliar diseases.

Jockey contains a red pigment so that treated grain can be easily identified.

### Fungicide Resistance Warning

Jockey Systemic Seed Fungicide is a member of the triazole group of fungicides. For fungicide resistance management Jockey is a Group **C** fungicide.

Some naturally occurring individual fungi resistant to Jockey and other Group **C** fungicides may exist through normal genetic variability in any fungal population. The resistant individuals can eventually dominate the fungal population if these fungicides are used repeatedly. These resistant fungi will not be controlled by Jockey and other Group **C** fungicides, thus resulting in a reduction in efficacy and possible yield loss.

Since the occurrence of resistant fungi is difficult to detect prior to use, Bayer CropScience Pty Ltd accepts no liability for any losses that may result from the failure of Jockey to control resistant fungi.

### Export of Treated Produce

Growers should note that suitable MRLs or import tolerances may not be established in all markets for produce treated with Jockey Systemic Seed Fungicide. If you are growing produce for export, please check with Bayer CropScience Pty Ltd for the latest information on MRLs and import tolerances BEFORE using Jockey Systemic Seed Fungicide.

### Application

Jockey is applied to seed prior to sowing. Application must be conducted with a seed treatment machine suitable for flowable seed dressings, with an adjustable output. The equipment must be calibrated for the flow rate of the seed, to ensure even coverage and correct application rates.

For best results, Jockey should be used to treat only undamaged seed of high viability.

#### **Wheat**

Jockey is a water based flowable product that can be applied neat or diluted with water. When applying the diluted product, prepare sufficient solution for the quantity of grain to be treated. Do not store diluted product. Do not dilute Jockey with more than 1.5 litres of water for each tonne of seed treated.

#### **Canola**

Apply Jockey neat to canola seed. Do not add water.

### Mixing Instructions

#### **Wheat only**

If mixing other compatible products or water with Jockey, mix only sufficient product for immediate use. Add the required compatible product or water to Jockey and agitate thoroughly by shaking, rolling or stirring. Do not allow mixed product to stand for more than five hours without further agitation.

**Compatibility****Wheat only**

Jockey is compatible with Vincit<sup>®</sup> C, Vitaflo<sup>®</sup> C (up to 1.25 L/tonne) and Raxil<sup>®</sup> C seed treatments.

Where additional protection against grain storage pests is required, Bayer CropScience Pty Ltd recommends a mixture of Jockey with K-Obiol<sup>®</sup> or another registered grain protection insecticide. Refer to the product label for application information.

**Crop Safety****Canola**

Jockey may shorten the hypocotyl length of canola. To avoid possible negative effects, Bayer CropScience recommends following good canola sowing practices. Avoid sowing Jockey treated canola deeper than 20 mm, or into soils prone to crusting. Ensure Jockey treated seed is sown in the season of treatment.

**PRECAUTIONS**

DO NOT use treated seed for human or animal consumption. DO NOT allow treated seed to contaminate grain or other seed intended for human or animal consumption.

**PROTECTION OF WILDLIFE, FISH, CRUSTACEANS AND ENVIRONMENT**

Treated seed could be harmful to game and wildlife.

DO NOT feed treated seed to wild or domestic birds. Any spillage of treated seed, however minor, must be cleaned up immediately, preferably recovered and used.

DO NOT contaminate streams, rivers or waterways with either this product, washings of this product, treated seed, or any container which has held product (see Storage and Disposal for directions).

**PROTECTION OF LIVESTOCK**

DO NOT feed treated seed to animals, including poultry.

**STORAGE AND DISPOSAL****Product and original container**

Store in the closed, original container in a cool, well-ventilated area. Do not store for long periods in direct sunlight. Do not store near foodstuffs or animal feed.

**5, 10, 20 & 200 L containers**

Triple or preferably pressure rinse containers before disposal. Dispose of rinsings in disposal pit, or use for diluting product to required volume. Do not dispose of undiluted chemicals on site. If recycling, replace cap and return clean containers to recycler or designated collection point. If not recycling, break, crush, or puncture and bury empty containers in a local authority landfill. If no landfill is available, bury the containers below 500 mm in a disposal pit specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots. Empty containers and product should not be burnt.

**100 and 110 L containers**

Empty container by pumping through the dry-break connection system. Do not attempt to unscrew the valve or breach the locked filling point. Do not contaminate the container with water or other foreign material.

Ensure that the coupler, pump, meter and hoses are disconnected, triple rinsed with clean water and drained after each use. When empty, or contents no longer required, return the container to the point of purchase. This container remains the property of Bayer CropScience Pty Ltd.

**Application equipment**

Rinse all application equipment with clean water immediately after use and dispose of rinsings in a dedicated disposal pit, specifically marked and set up for this purpose clear of waterways, desirable vegetation and tree roots.

**Treated seed and containers of treated seed**

Store treated seed in clearly marked bags or silos, away from other grain, animal feed or foodstuffs, in a cool, dry storage area. Do not use treated seed for human or animal consumption. If disposal of treated seed is necessary e.g. when cleaning up spillages, bury seed below 500 mm in a suitably located disposal pit, as described above. Do not use bags that have contained treated seed for any other purpose.

**SAFETY DIRECTIONS**

Harmful if swallowed. May irritate the eyes. Avoid contact with the eyes. When preparing the mixture and using the prepared mixture, wear cotton overalls buttoned to the neck and wrist (or equivalent clothing), elbow-length PVC gloves and goggles. Wash hands after use. After each day's use, wash gloves, goggles and contaminated clothing.

**FIRST AID**

If poisoning occurs contact a doctor or Poisons Information Centre (telephone: 13 11 26).

**MATERIAL SAFETY DATA SHEET**

Additional information is listed in the Material Safety Data Sheet, which can be obtained from [www.bayercropscience.com.au](http://www.bayercropscience.com.au).

**EXCLUSION OF LIABILITY**

This product must be used strictly as directed, and in accordance with all instructions appearing on the label and in other reference material. So far as it is lawfully able to do so, Bayer CropScience Pty Ltd accepts no liability or responsibility for loss or damage arising from failure to follow such directions and instructions.

Jockey®, Raxil® C and K-Obiol® are Registered Trademarks of Bayer.

NRA Approval No.: 52622/0303



<b>PESTICIDE, LIQUID, TOXIC, N.O.S</b> <b>(contains fluquinconazole)</b>		
<b>UN No. 2902</b>	<b>PG III</b>	<b>HAZCHEM 2X</b>
FOR 24 HOUR SPECIALIST ADVICE IN EMERGENCY ONLY PHONE 1800 033 111		

**DIRECTIONS FOR USE**

**Restrains**

DO NOT apply to damaged, shot or pinched seed.

DO NOT apply treated seed from aircraft.

CROP	DISEASE	APPLICATION RATE	CRITICAL COMMENTS
Wheat	Take-all (Suppression) <b>(Gaeumannomyces graminis)</b>	<b>Undiluted</b> 4.5 L/ tonne of seed	Ensure even coverage of seed. Apply with an applicator designed for liquid seed treatments. Calibrate application equipment for the flow rate of the grain. Jockey may be diluted with water prior to application to improve coverage. If applying dilute, or in mixtures the maximum recommended application volume is 6 L per tonne of seed.  * Stripe Rust is controlled for up to 6 weeks after sowing, with good suppression thereafter.  ** Leaf Rust is controlled for up to 4 weeks after sowing, with good suppression thereafter
	Common Bunt <b>(Tilletia spp.)</b>	<b>Diluted</b> 4.5 L plus up to 1.5 L of water/ tonne of seed	
	Seed- and Soil-borne Flag Smut <b>(Urocystis agropyri)</b>		
	Loose Smut <b>(Ustilago tritici)</b>		
	Stripe Rust * <b>(Puccinia striiformis f. sp. tritici)</b>		
	Leaf Rust ** <b>(Puccinia recondita f. sp. tritici)</b>		
	Septoria tritici blotch (Suppression) <b>(Mycosphaerella graminicola)</b>		
Canola	Blackleg (Suppression) <b>(Leptosphaeria maculans)</b>	20 L/tonne of seed	Ensure even coverage of seed. Apply undiluted with an applicator designed for liquid seed treatments. Calibrate application equipment for the flow rate of the grain.

**NOT TO BE USED FOR ANY PURPOSE OR IN ANY MANNER CONTRARY TO THIS LABEL UNLESS AUTHORISED UNDER APPROPRIATE LEGISLATION**

**WITHHOLDING PERIODS**

**Wheat**

Harvest: NOT REQUIRED WHEN USED AS DIRECTED

Grazing: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED, OR CUT FOR STOCK FOOD, WITHIN 12 WEEKS OF SOWING.

**Canola**

Harvest: NOT REQUIRED WHEN USED AS DIRECTED

Grazing: DO NOT GRAZE PLANTS GROWN FROM TREATED SEED, OR CUT FOR STOCK FOOD WITHIN 8 WEEKS OF SOWING.

**PROTECTION STATEMENT**

**DO NOT FEED TREATED SEED TO LIVESTOCK, INCLUDING POULTRY.**