

Supplemental Labeling



Dow AgroSciences LLC

9330 Zionsville Road

Indianapolis, IN 46268-1054 USA

WideMatch™

EPA Reg. No. 62719-512

Revised Rotation Interval for Field Peas

(For Distribution and Use in All States Except California, Idaho, Nevada, Oregon, Utah and Washington)

ATTENTION

- It is a violation of Federal law to use this product in a manner inconsistent with its labeling.
- This labeling must be in the possession of the user at the time of application.
- Read the label affixed to the container for WideMatch™ herbicide before applying. Carefully follow all precautionary statements and applicable use directions.
- Use of WideMatch according to this supplemental labeling is subject to all use precautions and limitations imposed by the label affixed to the container for WideMatch.

Directions for Use

Crop Rotation Intervals

Note: Numbers in parenthesis and † refer to footnotes following tables.

Rotation Crops (1)	Rotation Interval † (Soils greater than 2% organic matter AND rainfall more than 15 inches during 12 months following application)	Rotation Interval † (Soils less than 2% organic matter AND rainfall less than 15 inches during 12 months following application)
barley, grasses, field corn, oats, sweet corn, wheat	Anytime	Anytime
canola (rapeseed), cole crops (<i>Brassica</i> species), flax, garden beet, popcorn, spinach, sugarbeet, turnip	120 days	120 days
alfalfa, asparagus, grain sorghum, mint, onions, safflower, strawberries	10.5 months	10.5 months
field peas (4), dry beans, soybeans, sunflowers	10.5 months	18 months (2)
chick peas, lentils, potatoes (including potatoes grown for seed), and broadleaf crops grown for seed (excluding <i>Brassica</i> species)	18 months (2)	18 months (2, 3)

1. A field bioassay is recommended prior to planting any broadleaf crops that are not listed. Do not rotate to unlisted crops prior to 10.5 months following application.
2. An 18-month crop rotation is recommended due to the potential for crop injury. **Note:** For these crops, a minimum 10.5 month rotation interval must be observed to avoid illegal residues in the harvested crop.
3. A field bioassay is also recommended prior to planting these sensitive crops. See instructions on label affixed to container for WideMatch.
4. For rotation to field peas in 10.5 months, precipitation must be greater than 7.0 inches during the 10.5 months following application of WideMatch and greater than 5.5 inches during the June 1 through August 31 time period following application. Otherwise, rotation to field peas is recommended 18 months following application.

† **Note:** The above crop rotation intervals are based on average annual precipitation, regardless of irrigation practices. Observance of recommended crop rotation intervals should result in adequate safety to rotational crops. However, WideMatch is dissipated in the soil by microbial activity and the rate of microbial activity is dependent on several interrelating factors including soil moisture, temperature and organic matter. Therefore, accurate prediction of rotational crop safety is not possible. In areas of low organic matter (<2.0%) and less than 15 inches average annual precipitation, potential for crop injury may be reduced by burning or removal of plant residues, supplemental fall irrigation and deep moldboard plowing prior to planting the sensitive crop.

™Trademark of Dow AgroSciences LLC

R202-001

Accepted 05/06/05

Initial printing.