



# Predict<sup>®</sup>

## Herbicide

For control of grass and broadleaf weeds in field-grown nursery stock and noncrop land

*Active Ingredient:*

Norflurazon:  
4-chloro-5-(methylamino)-  
2-( $\alpha,\alpha,\alpha$ -trifluoro-*m*-tolyl)-  
3-(2*H*)-pyridazinone . . . . . 78.6%\*

*Other Ingredients:* 21.4%

*Total:* 100.0%

*\*Based on technical active ingredient analysis by isomer specific method AM-0864. Previously 80% by method T-4295.*

**SCP 849B-L1E 0201  
155228**

**10 pounds**  
Net Weight

**KEEP OUT OF REACH OF CHILDREN.**

## CAUTION

See additional precautionary statements on back of bag.

EPA Reg. No. 100-849

EPA Est. 55618-SC-001



## CONDITIONS OF SALE AND LIMITATION OF WARRANTY AND LIABILITY

**NOTICE:** Read the entire Directions for Use and Conditions of Sale and Limitation of Warranty and Liability before buying or using this product. If the terms are not acceptable, return the product at once, unopened, and the purchase price will be refunded.

The Directions for Use of this product should be followed carefully. It is impossible to eliminate all risks inherently associated with the use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as manner of use or application, weather or crop conditions, presence of other materials or other influencing factors in the use of the product, which are beyond the control of SYNGENTA CROP PROTECTION, Inc. or Seller. All such risks shall be assumed by Buyer and User, and Buyer and User agree to hold SYNGENTA and Seller harmless for any claims relating to such factors.

SYNGENTA warrants that this product conforms to the chemical description on the label and is reasonably fit for the purposes stated in the Directions for Use, subject to the inherent risks referred to above, when used in accordance with directions under normal use conditions. This warranty does not extend to the use of the product contrary to label instructions, or under abnormal conditions or under conditions not reasonably foreseeable to or beyond the control of Seller or SYNGENTA, and Buyer and User assume the risk of any such use. SYNGENTA MAKES NO WARRANTIES OF MERCHANTABILITY OR OF FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

In no event shall SYNGENTA or Seller be liable for any incidental, consequential or special damages resulting from the use or handling of this product. **THE EXCLUSIVE REMEDY OF THE USER OR BUYER, AND THE EXCLUSIVE LIABILITY OF SYNGENTA AND SELLER FOR ANY AND ALL CLAIMS, LOSSES, INJURIES OR DAMAGES (INCLUDING CLAIMS BASED ON BREACH OF WARRANTY, CONTRACT, NEGLIGENCE, TORT, STRICT LIABILITY OR OTHERWISE) RESULTING FROM THE USE OR HANDLING OF THIS PRODUCT, SHALL BE THE RETURN OF THE PURCHASE PRICE OF THE PRODUCT OR, AT THE ELECTION OF SYNGENTA OR SELLER, THE REPLACEMENT OF THE PRODUCT.**

SYNGENTA and Seller offer this product, and Buyer and User accept it, subject to the foregoing conditions of sale and limitations of warranty and of liability, which may not be modified except by written agreement signed by a duly authorized representative of SYNGENTA.

## GENERAL USE DIRECTIONS

It is a violation of Federal law to use this product in a manner inconsistent with its labeling.

Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation.

### AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for the protection of agricultural workers on farms, forests, nurseries, and greenhouses, and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification, and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE) and restricted-entry interval. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

**Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.** Exception: If the product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water is:

- Coveralls
- Waterproof gloves
- Shoes plus socks

Before applying Predict, read all directions and precautions appearing on the container label and this booklet.

**FAILURE TO FOLLOW THE DIRECTIONS FOR USE AND PRECAUTIONS ON THIS LABEL MAY RESULT IN POOR WEED CONTROL, CROP INJURY, OR ILLEGAL RESIDUES.**

**NOT FOR SALE, USE, OR DISTRIBUTION IN NASSAU COUNTY AND SUFFOLK COUNTY, NEW YORK.**

## GENERAL INFORMATION

Predict is a soil active herbicide for control of certain grasses and broadleaf weeds in field-grown nursery stock and noncrop lands.

Predict must be moved into the weed seed germination zone to be effective. If no rainfall occurs within 4 weeks after application, the product must be activated by flood irrigation or sprinkler irrigation. Predict alone has no postemergence activity and will not control established weeds. Existing weeds must be mechanically removed or controlled by using a suitable postemergence herbicide.

Cover crops planted in treated areas must not be harvested, grazed, or fed to livestock.

# Predict®

**Precautions for Field-Grown Nursery Stock:** (1) To avoid plant injury, do not apply until the fall following the first full season of field growth after transplanting. (2) Use only on recommended plantings (see following list) grown on medium- and fine-textured soils. **Do not use on coarse-textured soils.** (3) Caution should be taken when application is made to field-grown plants when roots are planted bound (i.e., in synthetic or burlap bags, etc.), so that herbicide does not channel directly into the root zone. (4) Wait at least 12 months before planting cotton or crops listed on this label. Other crops should not be planted in previously treated soil until a bioassay or test planting with the desired crop exhibits growth without chlorosis for 4 months after emergence. (5) Do not apply to erodible soils which may wash into the root zone of sensitive plants.

## DIRECTIONS FOR FIELD-GROWN NURSERY STOCK APPLICATIONS

Apply Predict before weeds emerge. If weeds have germinated prior to the application of Predict, a contact or burndown herbicide must be used (see tank mix instructions below).

The soil should be settled and firm at the time of application.

Apply as a directed spray. Avoid contact with foliage. Make only one application per year.

Do not apply this product through any type of irrigation system, or contaminate irrigation ditches or water used for domestic purposes.

Apply the recommended rate of Predict in 20-100 gals. of water per acre.

### Equipment

Predict should be applied using an accurately calibrated fixed boom sprayer. Filters with screen sizes of 50-mesh or larger should be used.

### Mixing Instructions

Clean and calibrate the sprayer. Add Predict to the spray tank already filled with the required volume of water. This will eliminate or minimize foaming. Maintain agitation while filling and spraying. If a bypass line is used, discharge at the bottom of the tank to further minimize foaming.

Do not allow Predict spray mixture to remain in the spray tank overnight, because settling will occur and resuspension will be difficult.

### Tank Mix Recommendations for Field-Grown Nursery Stock

Tank mix combinations may be desired for broader spectrum preemergence control or postemergence control of emerged weeds. Predict may be tank mixed with other EPA-registered herbicides in accordance with all directions, precautions, limitations, and recommendations on the label of the tank mix partner, after a compatibility test has shown that no formulation compatibility problem(s) exist (see **Compatibility Test** section).

**Note:** Predict may not always be compatible with formulations of DNA herbicides.

Products should be added to the spray tank in the following order:

1. Wettable powders and water-dispersible granules: wettable powders should be premixed in a small amount of water; water-dispersible granules should be added during filling. Allow both to disperse before other products are added. The order of addition is not important.
2. Flowable liquids
3. Emulsifiable concentrates
4. Surfactants: Begin adding wettable powders, flowable liquids, emulsifiable concentrates, and surfactants after the spray tank is  $\frac{3}{4}$  full. Continue agitation during the addition of all the materials and while filling and spraying.

### Compatibility Test

Before mixing in the spray tank, it is advisable to test compatibility by mixing all components in a small container in proportionate quantities (see following table).

Herbicide Formulations	Amount of Herbicide to Add to One Pt. of Spray Carrier (Assuming Volume is 25 Gals. per Acre)	
	Rate Per Acre	Level Teaspoons
Dry	1 lb.	1.5
Liquid	1 pt.	0.5

If herbicide(s) do not ball-up or form flakes, sludge, gels, oily films or layers, or other precipitates, then the tested spray mix is compatible. Usually, incompatibility in any of the above described forms will occur within 5 minutes after mixing.

If components are incompatible, the use of a compatibility agent approved for agricultural use is recommended. Rerun the above compatibility test with a suitable compatibility agent ( $\frac{1}{4}$  teaspoon is equivalent to 2 pts./100 gals. of spray carrier).

### Broadcast Rates

Apply 3 lbs. of product per acre in the fall or spring.

### Strip or Band Treatment

When applying a strip treatment of Predict, the following formula may be used to calculate the amount per acre:

$$\frac{\text{Width of strip in ft.}}{\text{Distance between rows in ft.}} \times \text{Lbs. per acre for broadcast treatment} = \text{Lbs. per acre for strip treatment}$$

The solution should be mixed to the maximum label rate and at no point in the field should the solution be applied as a concentration any higher than this rate.

## RECOMMENDATIONS FOR FIELD-GROWN NURSERY STOCK CROPS

Apply Predict to these species of trees and shrubs as recommended above:

Apple	<i>Malus</i>
Blueberry	<i>Vaccinium</i>
Citrus	<i>Citrus</i>
Crabapple*	<i>Malus*</i>
Cranberry*	<i>Viburnum*</i>
Filbert	<i>Corylus</i>
Pear	<i>Pyrus</i>
Pecan	<i>Carya</i>
Walnut	<i>Juglans</i>
Boxwood, Korean	<i>Buxus harlandii</i>
Cypress, Bald	<i>Taxodium distichum</i>
Dogwood	<i>Cornus</i>
Holly,	
Green Luster	<i>Ilex crenata</i> 'Green Luster'
East Palaka	<i>I. opaca</i> 'East Palaka'
American Greenleaf	<i>I. X</i> 'American Greenleaf'
Foster	<i>I. X</i> 'Fosteri'
Nellie Stevens	<i>I. X</i> 'Nellie Stevens'
Savannah	<i>I. X</i> 'Savannah'
Juniper,	
Andora Creeping	<i>Juniperus horizontalis</i> 'Plumosa'
Burki	<i>J. virginiana</i> 'Burki'
Fastigata	<i>J. fastigata</i>
Magnolia, Southern	<i>Magnolia grandiflora</i>
Maple, Red	<i>Acer rubrum</i>
Myrtle, Crepe	<i>Lagerstroemia indica</i>
Oak,	
Pin	<i>Quercus palustris</i>
Willow	<i>Q. phellos</i>
Poplar, Tulip	<i>Liriodendron tulipifera</i>
Sycamore, American	<i>Platanus occidentalis</i>

\*Fruit is not to be used for feed and/or human consumption.

## WEEDS CONTROLLED AT FIELD-GROWN NURSERY STOCK RATES

Predict at recommended rates controls the following weeds:

### Broadleaf Weeds (Dicotyledons)

Carolina Geranium (wild)	<i>Geranium carolinianum</i>
Carpetweed	<i>Mullugo verticillata</i>
Common Chickweed	<i>Stellaria media</i>
Desert Rockpurslane (Redmaids)	<i>Calandrinia ciliata</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Falsedandelion (Smooth Cat's Ear)	<i>Pyrrhopappus carolinianus</i>
Fiddleneck	<i>Amsinckia intermedia</i>
Filaree* (Redstem and Whitestem)	<i>Erodium</i> spp.*
Little Mallow (Cheeseweed)	<i>Malva parviflora</i>
London Rocket	<i>Sisymbrium irio</i>
Pigweeds	<i>Amaranthus</i> spp.
Pineappleweed	<i>Matricaria matricarioides</i>
Prickly Sida	<i>Sida spinosa</i>
Prostrate Spurge	<i>Euphorbia humistrata</i>
Puncturevine	<i>Tribulus terrestris</i>
Purslane	<i>Portulaca oleracea</i>
Purple Cudweed	<i>Gnaphalium purpureum</i>
Shepherdspurse	<i>Capsella bursa-pastoris</i>
Spurred Anoda	<i>Anoda cristata</i>
Stinging Nettle	<i>Urtica dioica</i>
Tropic Croton	<i>Croton glandulosus</i>
Tumble Mustard	<i>Sisymbrium altissimum</i>
Velvetleaf	<i>Abutilon theophrasti</i>
Wild Buckwheat	<i>Polygonum convolvulus</i>

\*Treat prior to germination and incorporate with water on medium soils if necessary for adequate control.

## Predict®

### Grass and Sedge Weeds (Monocotyledons)

Annual Bluegrass	<i>Poa annua</i>
Annual Sedge	<i>Cyperus compressus</i>
Bahiagrass (seedling)	<i>Paspalum notatum</i>
Barlyardgrass	<i>Echinochloa crus-galli</i>
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>
Brome	<i>Bromus</i> spp.
Cheat	<i>Bromus secalinus</i>
Crabgrass	<i>Digitaria</i> spp.
Downy Brome	<i>Bromus tectorum</i>
Fall Panicum	<i>Panicum dichotomiflorum</i>
Feather Fingergrass	<i>Chloris virgata</i>
Foxtails	<i>Setaria</i> spp.
Goosegrass	<i>Eleusine indica</i>
Italian Ryegrass (Annual Ryegrass)	<i>Lolium multiflorum</i>
Johnsongrass (seedling)	<i>Sorghum halepense</i>
Sandbur	<i>Cenchrus longispinus</i>
Sixweeks Grama	<i>Bouteloua barbata</i>
Southwestern Cupgrass	<i>Eriochloa gracilis</i>
Tall Fescue	<i>Festuca arundinacea</i>
Texas Panicum	<i>Panicum texanum</i>
Wild Barley	<i>Hordeum leporinum</i>
Wild Onion	<i>Allium canadense</i>
Witchgrass	<i>Panicum capillare</i>

#### Predict applied at recommended rates suppresses the following grass and broadleaf weeds:

Bermudagrass	<i>Cynodon dactylon</i>
Camphorweed	<i>Heterotheca subaxillaris</i>
Cocklebur	<i>Xanthium strumarium</i>
Common Lambsquarters	<i>Chenopodium album</i>
Common Ragweed	<i>Ambrosia artemisiifolia</i>
Florida pusley	<i>Richardia scabra</i>
Goldenrod	<i>Solidago altissima</i>
Groundsel	<i>Senecio vulgaris</i>
Hairy Fleabane (Flax-leaved Fleabane)	<i>Conyza bonariensis</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed (Marestail)	<i>Conyza canadensis</i>
Johnsongrass (rhizome)	<i>Sorghum halepense</i>
Kochia	<i>Kochia scoparia</i>
Lanceleaf Sage	<i>Salvia retroflexa</i>
Morningglory	<i>Ipomoea</i> spp.
Natalgrass (seedling)	<i>Rhynchelytrum repens</i>
Nutsedge	<i>Cyperus</i> spp.
Orchardgrass	<i>Dactylis glomerata</i>
Pangolagrass (seedling)	<i>Digitaria decumbens</i>
Pennsylvania Smartweed	<i>Polygonum pensylvanicum</i>
Plantains (Bracted and Buckhorn)	<i>Plantago</i> spp.
Poorjoe	<i>Diodia teres</i>
Quackgrass	<i>Agropyron repens</i>
Russian thistle	<i>Salsola iberica</i>
Silverleaf Nightshade (White Horsenettle)	<i>Solanum elaeagnifolium</i>
Sowthistle	<i>Sonchus oleracea</i>
Spotted Spurge	<i>Euphorbia maculata</i>
Vaseygrass	<i>Paspalum urvillei</i>
Wirestem Muhly (Western Muhly)	<i>Muhlenbergia frondosa</i>

### DIRECTIONS FOR NONCROP LANDS

Predict may be used for preemergence weed control in noncrop land areas including: industrial sites, rights-of-way (highway, pipeline, railroad, or utility), wasteland, and other noncrop land areas. Do not apply to erodible soils which may wash into the root zone of sensitive plants.

Use Predict at a rate of 2.5-5 lbs. of product per treated acre for noncrop land areas. Higher rates within the range should be used for fine-textured soils and where longer residual is desired.

Apply the recommended rate of Predict in 20-100 gals. of water per acre.

Since Predict is a preemergence herbicide, it must be applied to the soil surface before weeds germinate. Existing weeds should be mechanically removed or controlled with a suitable postemergence herbicide. Predict must be incorporated into the soil by rainfall or irrigation within 4 weeks of application for best weed control.

#### Tank mix recommendations for noncrop lands

Tank mix combinations may be desired for broader spectrum preemergence control or postemergence control of emerged weeds or brush. Predict may be tank mixed with other EPA-registered herbicides in accordance with all directions, precautions, limitations, and recommendations on the label of the tank mix partner, after a compatibility test has shown that no formulation compatibility problem(s) exist (see **Compatibility Test** section).

**Note:** Predict may not always be compatible with formulations of DNA herbicides.

**WEEDS CONTROLLED AND SUPPRESSED IN NONCROP LANDS**

Predict at recommended rates controls the following weeds:

**Broadleaf Weeds (Dicotyledons)**

Annual Sedge	<i>Cyperus compressus</i>
Camphorweed	<i>Heterotheca subaxillaris</i>
Carolina Geranium (wild)	<i>Geranium carolinianum</i>
Common Chickweed	<i>Stellaria media</i>
Common Ragweed	<i>Ambrosia artemisiifolia</i>
Desert Rockpurslane (Redmaids)	<i>Calandrinia ciliata</i>
Dogfennel	<i>Eupatorium capillifolium</i>
Falsedandelion (Smooth Cat's Ear)	<i>Pyrrhopappus carolinianus</i>
Fiddleneck	<i>Amsinckia intermedia</i>
Filaree* (Redstem and Whitestem)	<i>Erodium</i> spp.*
Goldenrod	<i>Solidago altissima</i>
Little Mallow (Cheeseweed)	<i>Malva parviflora</i>
London Rocket	<i>Sisymbrium irio</i>
Pineappleweed	<i>Matricaria matricarioides</i>
Prostrate Spurge	<i>Euphorbia humistrata</i>
Puncturevine	<i>Tribulus terrestris</i>
Purple Cudweed	<i>Gnaphalium purpureum</i>
Shepherdspurge	<i>Capsella bursa-pastoris</i>
Spreading Dayflower	<i>Commelina diffusa</i>
Stinging Nettle	<i>Urtica dioica</i>
Tumble Mustard	<i>Sisymbrium altissimum</i>
Wild Buckwheat	<i>Polygonum convolvulus</i>

\*Treat prior to germination and incorporate with water on medium soils if necessary for adequate control.

**Grass and Sedge Weeds (Monocotyledons)**

Annual Bluegrass	<i>Poa annua</i>
Annual Sedge	<i>Cyperus compressus</i>
Bahiagrass (seedling)	<i>Paspalum notatum</i>
Barnyardgrass	<i>Echinochloa crus-galli</i>
Bearded Sprangletop	<i>Leptochloa fascicularis</i>
Broadleaf Signalgrass	<i>Brachiaria platyphylla</i>
Cheat	<i>Bromus secalinus</i>
Crabgrass	<i>Digitaria</i> spp.
Crowfootgrass (seedling)	<i>Dactyloctenium aegyptium</i>
Downy Brome	<i>Bromus tectorum</i>
Fall Panicum	<i>Panicum dichotomiflorum</i>
Feather Fingergrass	<i>Chloris virgata</i>
Foxtails	<i>Setaria</i> spp.
Goosegrass	<i>Eleusine indica</i>
Guineagrass (seedling)	<i>Panicum maximum</i>
Italian Ryegrass (Annual Ryegrass)	<i>Lolium multiflorum</i>
Johnsongrass (seedling)	<i>Sorghum halepense</i>
Natalgrass (seedling)	<i>Rhynchelytrum repens</i>
Pangolagrass (seedling)	<i>Digitaria decumbens</i>
Sandbur	<i>Cenchrus longispinus</i>
Sixweeks Grama	<i>Bouteloua barbata</i>
Southwestern Cupgrass	<i>Eriochloa gracilis</i>
Tall Fescue	<i>Festuca arundinacea</i>
Texas Panicum	<i>Panicum texanum</i>
Vaseygrass (seedling)	<i>Paspalum urvillei</i>
Wild Barley	<i>Hordeum leporinum</i>
Wild Onion	<i>Allium canadense</i>
Witchgrass	<i>Panicum capillare</i>

## Predict®

### Predict applied at recommended rates suppresses the following weeds:

Bermudagrass	<i>Cynodon dactylon</i>
Common Lambsquarters	<i>Chenopodium album</i>
Common Purslane	<i>Portulaca oleracea</i>
Florida Pusley	<i>Richardia scabra</i>
Groundsel	<i>Senecio vulgaris</i>
Hairy Fleabane (Flax-leaved Fleabane)	<i>Conyza bonariensis</i>
Henbit	<i>Lamium amplexicaule</i>
Horseweed (Marestail)	<i>Conyza canadensis</i>
Johnsongrass (rhizome)	<i>Sorghum halepense</i>
Nutsedge	<i>Cyperus</i> spp.
Orchardgrass	<i>Dactylis glomerata</i>
Pigweeds (Redroot, Tumble, and Green Amaranth)	<i>Amaranthus</i> spp.
Plantains (Bracted and Buckhorn)	<i>Plantago</i> spp.
Poorjoe	<i>Diodia teres</i>
Russian Thistle	<i>Salsola iberica</i>
Quackgrass	<i>Agropyron repens</i>
Silverleaf Nightshade	<i>Solanum elaeagnifolium</i>
Sowthistle	<i>Sonchus oleracea</i>
Torpedograss	<i>Panicum repens</i>
Wirestem Muhly (Western Muhly)	<i>Muhlenbergia frondosa</i>

## STORAGE AND DISPOSAL

### Storage

Do not contaminate water, food, or feed by storage or disposal.

### Pesticide Disposal

Pesticide wastes are toxic. Improper disposal of excess pesticide, spray mixture, or rinsate is a violation of Federal law. If waste cannot be disposed of by use of label instructions, contact your state pesticide or environmental control agency or hazardous waste representative at the nearest EPA Regional Office for guidance.

### Container Disposal

Completely empty container into application equipment. Then dispose of empty container in a sanitary landfill or by incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

For minor spills, leaks, etc., follow all precautions indicated on this label and clean up immediately. Take special care to avoid contamination of equipment and facilities during cleanup procedures and disposal of wastes. In the event of a major spill, fire, or other emergency, call 1-800-888-8372, day or night.

## PRECAUTIONARY STATEMENTS

### Hazards to Humans and Domestic Animals

#### CAUTION

Harmful if swallowed or absorbed through the skin. Avoid contact with skin, eyes, or clothing. In case of skin or eye contact, flush with plenty of water.

### Statement of Practical Treatment

**If swallowed:** Call a physician or poison control center. Drink 1 or 2 glasses of water and induce vomiting by touching back of throat with finger or if available by administering syrup of ipecac. Do not induce vomiting or give anything by mouth to an unconscious person.

**If inhaled:** Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

**If on skin:** Wash with plenty of soap and water. Get medical attention if irritation persists.

### Personal Protective Equipment

#### Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Waterproof gloves
- Shoes plus socks

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

### Engineering Control Statements

When handlers use closed systems, enclosed cabs, or aircraft in a manner that meets the requirements listed in the Worker Protection Standard (WPS) for agricultural pesticides [40 CFR 170.240(d)(4-6)], the handler PPE requirements may be reduced or modified as specified in the WPS.

*continued...*

**PRECAUTIONARY STATEMENTS (continued)**

**User Safety Recommendations**

**Users should:**

- Wash hands before eating, drinking, chewing gum, using tobacco, or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.
- Remove PPE immediately after handling this product. Wash the outside of gloves before removing. As soon as possible, wash thoroughly and change into clean clothing.

**Environmental Hazards**

Do not contaminate water when disposing of equipment wash water. Do not apply directly to water, or to areas where surface water is present, or to intertidal areas below the mean high water mark. Do not allow this material to drift onto neighboring crops or noncrop areas or use in a manner or at a time other than in accordance with label directions because animal, plant or crop injury, illegal residues, or other undesirable results may occur.

Norflurazon can contaminate surface water through spray drift. Under some conditions, norflurazon may also have a high potential for runoff into surface water (primarily via dissolution in runoff water) for several months post-application. These include poorly draining or wet soils with readily visible slopes toward adjacent surface waters, frequently flooded areas, areas overlaying extremely shallow ground water, areas with in-field canals or ditches that drain to surface water, areas not separated from adjacent surface waters with vegetated filter strips, and areas overlaying tile drainage systems that drain to surface water.

This chemical is known to leach through soil into ground water under certain conditions as a result of agricultural use. Use of this chemical in areas where soils are permeable, particularly where the water table is shallow, may result in ground water contamination.

Predict® trademark of Syngenta

©2001 Syngenta

Syngenta Crop Protection, Inc.  
Greensboro, North Carolina 27409  
www.syngenta-us.com

**SCP 849B-L1E 0201**  
**155228**