



Bayer CropScience LP
 P.O. Box 12014
 2 T.W. Alexander Drive
 Research Triangle Park, North Carolina 27709
 1-866-99BAYER (1-866-992-2937)
<http://www.bayercropscienceus.com>

Flint® Fungicide
 EPA Reg. No. 264-777

For Use on Tree Nuts and Stone Fruits

PRODUCT BULLETIN

FOR DISTRIBUTION AND USE ONLY IN THE STATE OF CALIFORNIA

FIFRA Section 2(ee) Recommendation: This recommendation is made as permitted under FIFRA Section 2(ee) and has not been submitted to or accepted by the U.S. Environmental Protection Agency. It is a violation of federal law to use this product in a manner inconsistent with its label. This product bulletin must be in the possession of the user at the time of pesticide application. Follow all applicable directions, restrictions, Worker Protection Standard requirements, and precautions on the EPA registered label.

DIRECTIONS FOR USE

It is a violation of Federal law to use this product inconsistent with its labeling.

STONE FRUITS: Apricots, Cherries, Nectarines, Peaches, Plums, Plumcots, Prunes (fresh)			
Disease Control	Rate oz/Acre	Application Timing	Notes
Cherry Leaf Spot (<i>Blumeriella jaapii</i>) Powdery Mildew (<i>Podosphaera</i> spp. and <i>Sphaerotheca pannosa</i>) Scab (<i>Cladosporium carpophilum</i>) Rust (<i>Tranzschelia discolor</i>)	2.0-4.0	Begin applications preventively. Apply at petal fall and continue on a 7- to 14-day interval.	Use the higher rates and shorter intervals when disease pressure is severe.
Restrictions: Do not apply more than 16 oz of Flint® per acre per season. Do not apply Flint® within 1 day of harvest. Do not apply more than 4 applications of Flint® per acre per season.			

TREE NUTS: Almonds,* Beechnuts, Brazil Nuts, Butternuts, Cashew, Chestnuts, Chinquapins, Filberts, Hickory Nuts, Macadamia Nuts, Pecans,* Pistachios,* Walnuts (* See Specific Use Instructions)			
Disease Control	Rate Oz/Acre	Application Timing	Notes
Botryosphaeria Panicle and Shoot Blight (<i>Botryosphaeria dothidea</i>)	2-3	Begin applications preventively and continue as needed on a 14- to 21-day interval	Use the higher rates and shorter intervals when disease pressure is severe.
Alternaria Late Blight (<i>Alternaria alternata</i>) Anthracnose (<i>Colletotrichum acutatum</i> , <i>Glomerella cingulata</i>) Scab (<i>Cladosporium carpophilum</i> , <i>Cladosporium caryigenum</i>) Shothole (<i>Wilsonomyces carpophilus</i>) Rust (<i>Tranzschelia discolor</i>)	3-4	Begin applications preventively and continue as needed on a 7- to 14-day interval.	
Restrictions: Do not apply more than 16 oz of Flint® per acre per season. Do not apply Flint® within 60 days of harvest. Do not apply more than 4 applications of Flint® (or other strobilurin fungicides) alone or 4 applications of Flint® (or other strobilurin fungicides) in mixture with non-strobilurin fungicides.			

For MEDICAL And TRANSPORTATION Emergencies ONLY Call 24 Hours A Day 1-800-334-7577

For PRODUCT USE Information Call 1-866-99BAYER (1-866-992-2937)

Visit our worldwide web site at www.bayercropscienceus.com

As with any crop-protection product, always read and follow label directions.

For additional information call toll-free 1-866-99BAYER (1-866-992-2937).