



Trilin[®]

HERBICIDE

A Selective Herbicide for the Preemergence Control of Annual Grasses and Broadleaf Weeds

ACTIVE INGREDIENT

Trifluralin (a, a, a-trifluoro-2, 6-dinitro-N, N-dipropyl-p-toluidine)..... 42.8%

INERT INGREDIENTS 57.2%

TOTAL.....100.0%

Contains 4 Pounds Trifluralin Per Gallon

Contains Petroleum Distillate

**KEEP OUT OF REACH OF CHILDREN
WARNING – AVISO**

Si usted no entiende la etiqueta, busque a alguien para que se la explique a usted en detalle. (If you do not understand the label, find someone to explain it to you in detail.)

STATEMENT OF PRACTICAL TREATMENT

IF SWALLOWED: Drink 1 to 2 glasses of water. Induce vomiting by placing finger in back of throat. Call a physician. Do not induce vomiting or give anything by mouth to an unconscious person.

IF INHALED: Remove victim to fresh air. If not breathing, give artificial respiration, preferably mouth-to-mouth. Get medical attention.

IF ON SKIN: Wash immediately with soap and water.

IF IN EYES: Flush with plenty of water for at least 15 minutes. Get medical attention.

For medical emergencies involving this product, call toll free 1-800-237-1854.

See Label for Additional Precautions and Directions for Use

GRIFFIN L.L.C.

VALDOSTA, GEORGIA 31601

Specimen Label

EPA REG. NO. 1812-355

PRECAUTIONARY STATEMENTS HAZARDS TO HUMANS (AND DOMESTIC ANIMALS) WARNING

Causes substantial, but temporary eye injury. Do not get in eyes or on clothing. Harmful if swallowed, inhaled or absorbed through the skin. Avoid contact with skin or clothing. Do not contaminate foodstuffs or feeds.

PERSONAL PROTECTIVE EQUIPMENT

Some materials that are chemical-resistant to this product are listed below. If you want more options, follow the instructions for category G on an EPA chemical resistance category selection chart.

Applicators and other handlers must wear:

- Long-sleeved shirt and long pants
- Chemical-resistant gloves, such as barrier laminate, or viton \geq 14 mils
- Shoes plus socks
- Protective eyewear

Discard clothing and other absorbent materials that have been drenched or heavily contaminated with this product's concentrate. Do not reuse them.

Follow manufacturer's instructions for cleaning/maintaining PPE. If no such instructions for washables, use detergent and hot water. Keep and wash PPE separately from other laundry.

USER SAFETY RECOMMENDATIONS

Users should:

- Wash hands before eating, drinking, chewing gum, using tobacco or using the toilet.
- Remove clothing immediately if pesticide gets inside. Then wash thoroughly and put on clean clothing.

ENVIRONMENTAL HAZARDS

This pesticide is toxic to fish. Do not apply directly to water, to areas where surface water is present or to intertidal areas below the mean high water mark. Drift or runoff from treatment areas may be hazardous to aquatic organisms in neighboring aquatic sites. Do not contaminate water when disposing of equipment washwaters.

PHYSICAL AND CHEMICAL HAZARDS

Flammable. Keep away from heat and open flame.

DIRECTIONS FOR USE

It is a violation of Federal Law to use this product in a manner inconsistent with its labeling. Do not apply this product in a way that will contact workers or other persons, either directly or through drift. Only protected handlers may be in the area during application. For any requirements specific to your State or Tribe, consult the agency responsible for pesticide regulation. Trilin is a preemergence herbicide which is incorporated into the soil to provide control of grasses and broadleaf weeds. Trilin controls weeds as they germinate. Trilin will not control established weeds.

AGRICULTURAL USE REQUIREMENTS

Use this product only in accordance with its labeling and with the Worker Protection Standard, 40 CFR part 170. This Standard contains requirements for protection of agricultural workers on farms, forests, nurseries, and greenhouses and handlers of agricultural pesticides. It contains requirements for training, decontamination, notification and emergency assistance. It also contains specific instructions and exceptions pertaining to the statements on this label about personal protective equipment (PPE), and restricted-entry intervals. The requirements in this box only apply to uses of this product that are covered by the Worker Protection Standard.

Do not enter or allow worker entry into treated areas during the restricted-entry interval (REI) of 12 hours.

Exception: If product is soil-injected or soil-incorporated, the Worker Protection Standard, under certain circumstances, allows workers to enter the treated area if there will be no contact with anything that has been treated.

PPE required for early entry to treated areas that is permitted under the Worker Protection Standard and that involves contact with anything that has been treated, such as plants, soil, or water, is:

- Coveralls
- Chemical-resistant gloves, such as barrier laminate, or viton \geq 14 mils
- Shoes plus socks
- Protective eyewear

STORAGE AND DISPOSAL

Avoid freezing. Store above 40°F. If frozen, poor weed control may result. Do not store near heat or flame. Do not contaminate water, food or feed by storage or disposal.

PESTICIDE DISPOSAL: Wastes resulting from the use of this product may be disposed of on site or at an approved waste disposal facility.

CONTAINER DISPOSAL: Triple rinse (or equivalent). Then offer for recycling or reconditioning or puncture and dispose of in a sanitary landfill, or incineration, or, if allowed by state and local authorities, by burning. If burned, stay out of smoke.

GRASSES AND BROADLEAF WEEDS CONTROLLED BY TRILIN

Grasses

Annual Bluegrass	<i>Poa annua</i>
Barnyardgrass (Watergrass)	<i>Echinochloa</i> sp.
Brachiaria (Signalgrass)	<i>Brachiaria</i> sp.
Bromegrass (Cheatgrass) (Downy Brome)	<i>Bromus tectorum</i>
Cheat (Chess)	<i>Bromus secalinus</i>
Crabgrass (Large Crabgrass) (Smooth Crabgrass)	<i>Digitaria</i> spp.
Foxtail (Bottlegrass) (Bristlegrass) (Giant Foxtail) (Green Foxtail) (Foxtail Millet) (Pigeongrass) (Robust Foxtail) (Yellow Foxtail)	<i>Setaria</i> spp.
Itchgrass (Raoulgrass) (See Sugarcane for Special Instructions)	<i>Rottboellia exaltata</i>

Broadleaf Weeds

Carpetweed	<i>Mollugo verticillata</i>
Chickweed	<i>Stellaria media</i>
Field Bindweed (See Fruit and Nut Crops and Vineyards for Special Instructions)	<i>Convolvulus arvensis</i>
Florida Pusley (Florida Purslane) (Mexican Clover) (Pusley)	<i>Richardia scabra</i>
Goosefoot	<i>Chenopodium hybridum</i>
Henbit (Fall Application Only)	<i>Lamium amplexicaule</i>
Knotweed	<i>Polygonum aviculare</i>
Kochia (Fireweed) (Mexican Fireweed)	<i>Kochia scoparia</i>

Johnsongrass (from seed) (Rhizome – See Special Instructions for Control in Cotton, Soybeans, Fruit and Nut Crops, Vineyards and Cottonwood Trees Grown for Pulpwood)	<i>Sorghum halepense</i>
Junglerice	<i>Echinochloa colonum</i>
Panicum Fall Panicum (Spreading Panicgrass – See Special Instructions in Cotton and Soybeans)	<i>Panicum dichotomiflorum</i>
Guineagrass (See Sugarcane for Special Instructions)	<i>Panicum maximum</i>
Texas Panicum (See Special Instructions in Cotton and Soybeans) (Buffalograss) (Coloradograss)	<i>Panicum texanum</i>
Red Rice (See Suppression or Partial Control Directions Under Soybeans)	<i>Oryza sativa</i>
Sandbur (Burggrass)	<i>Cenchrus incertus</i>
Sprangletop	<i>Leptochloa filiformis</i>
Stinkgrass (Lovegrass)	<i>Eragrostis ciliaris</i>
Wild Cane (Shattercane) (See soybean – Trilin Alone for Special Instructions)	<i>Sorghum bicolor</i>
Woolly Cupgrass	<i>Eriochloa villosa</i>

Lambsquarters	<i>Chenopodium album</i>
Pigweed (Carelessweed) (Prostrate Pigweed) (Redroot) (Rough Pigweed) (Spiny Pigweed)	<i>Amaranthus</i> spp.
Puncturevine (Western U.S. Only) (Caltrop) (Goathead)	<i>Tribulus terrestris</i>
Purslane	<i>Portulaca oleracea</i>
Russian Thistle (Tumbleweed)	<i>Salsola kali</i>
Stinging Nettle (Nettle)	<i>Urtica dioica</i>

Long term and continued use of trifluralin has resulted in the selection of tolerant populations in certain species of weeds. This situation is limited to a few weeds and is generally geographically specific. Weed species known to have some trifluralin tolerant population are goosegrass, green foxtail (pigeongrass) and Palmer amaranthus (Palmer pigweed). Trilin is not recommended for the control of goosegrass, tolerant green foxtail or Palmer amaranthus. Consult State Agricultural Extension Service or Experiment Station weed specialist for specific recommendations for local weed populations.

SOIL PREPARATION

Soil surface should be smooth enough to enable the efficient use of sprayer and incorporation equipment to insure a uniform application and incorporation of Trilin. Interference can be caused by ground cover such as crop residue or existing weeds. Soil should be tilled prior to the application of Trilin to allow uniform incorporation into the top 2 to 3 inches of soil. Soil moisture should be such that any large clods will be broken up during incorporation process.

Crop Residues or Existing Weeds

Ground cover such as crop residues of existing weeds can interfere with the incorporation of Trilin into the soil. A manageable level of such ground cover will allow the Trilin to be uniformly incorporated into the top 2 to 3 inches of soil. If the level of the ground cover is such that this cannot be done you must till the soil prior to the application of Trilin.

Roughness

The soil surface should be smooth enough so that you can operate a sprayer and incorporation equipment efficiently and at speeds which insure a uniform application and incorporation of Trilin.

General Soil Conditions

To assure incorporation of Trilin, soil moisture conditions should be such that any large clods can be broken up during the incorporation process.

SOIL TEXTURE GUIDE

The amount of chemical applied will vary with the soil texture and organic matter. A fine textured soil will require more Trilin per acre than a coarse soil. Where rates are based on coarse, medium or fine textured soils, it is understood that soil textural classes are generally categorized as follows:

Coarse	Medium	Fine
Sand	Loam	Sandy clay
Loamy sand	Silt	Clay loam
Sandy loam	Silt loam	Silty clay, Clay
	Silty clay loam*	
	Sandy clay loam*	

*Silty clay loam and sandy clay loam soils are transitional soils and may be classified as either medium or fine textured soils. If silty clay loam or sandy clay loam soils are mostly sand or silt, they are usually classified as medium textured soils. If they are mostly clay, they are usually classified as fine textured soils.

Do not exceed recommended rates.

INCORPORATION DIRECTIONS

General Directions

Erratic weed control and/or crop injury may result if Trilin is not incorporated into the top 2 to 3 inches of the final seedbed.

Before Planting

Initial incorporation of Trilin must take place within 24 hours after application. A second incorporation prior to planting should be done by running the equipment in a different direction from the first. Incorporate Trilin uniformly into the top 2 to 3 inches of the final seedbed.

After Planting

For directions concerning incorporation after planting check label for crop or crops of interest for specific instructions.

72 Hour Incorporation Delay - Arizona and California Only

When Trilin is applied as a preplant incorporated treatment, the first incorporation pass must be accomplished within 24 hours after application.

In Arizona and California the incorporation delay has been extended from 24 to 72 hours when applied to dry soils. When Trilin is applied to warm soil or if wind velocity is 10 mph or higher, variable weed control may result from delaying the first incorporation beyond 24 hours. Where two incorporation passes are required, the second incorporation may occur anytime prior to planting.

Bedded Culture

Trilin needs to be incorporated into the top 2 to 3 inches of the final seedbed for effective weed control.

Application Prior To Bedding

Apply Trilin prior to bedding and incorporate with recommended equipment. The bedding operation serves as the second incorporation. Avoid removal of untreated soil from the seedbed before or during the planting operation. This would expose untreated soil allowing weeds to germinate in the drill row.

Application After Bedding

Knock off beds to planting height before application and incorporate Trilin with recommended equipment that will conform to the bed shape. Do not leave untreated soil exposed.

Recommended Equipment

Two incorporation passes are necessary unless specifically stated otherwise on this label. The second incorporation should not be deeper than the first.

Disc: Set disc to cut 4 to 6 inches deep and operate at 4 to 6 mph. A 4 to 6 inch cut should result in Trilin being incorporated into the top 2 to 3 inches of the soil.

Field Cultivator: Field cultivators are defined as implements with sweeps of 3 to 4 rows spaced at intervals of 7 inches or less staggered so that no soil is left unturned. Set to cut 3 to 4 inches deep and operate at 5 mph or more. Do not use chisel points.

Combination Seedbed Conditioners: Set to cut 3 to 4 inches deep and operate at a speed of at least 5 mph. These are defined as three or more tillage devices combined and used as a single tool. For example, two to three rows of field cultivator C- or S- shaped shanks with an effective sweep spacing of 6 to 9 inches (staggered so that no soil is left unturned) followed by a spike tooth or flexline harrow followed by a ground driven reel or basket.

Rolling Cultivator: Set to cut 2 to 4 inches deep and operate at a speed of 6 to 8 mph. Rolling cultivators are effective for use on coarse and medium textured soils. The rolling cultivator may be used on fine textured soils when used in sugarcane.

Bed Conditioner (Do-All): Set to cut 2 to 4 inches deep and operate at a speed of 4 to 6 mph. The Do-All is effective when used on coarse or medium textured soils only. Only one incorporation pass is necessary in bedded culture. Two passes with the Do-All are necessary in flat planted culture.

Mulch Treader (other similar disc-type implements): Set Mulch Treader to cut 3 to 4 inches deep and operate at 5 to 8 mph.

P.T.O. Driven Equipment (tillers, cultivators, hoes): Only one incorporation is required. Adjust to incorporate Trilin into the top 2 to 3 inches of the seedbed using rotors spaced to give a clean sweep of the soil. P.T.O. equipment should not be operated at a speed greater than 4 mph.

Other equipment, including the flexible tine-tooth harrow (Flexline, Melroe), is also recommended but only for the special usages for which it is specified in this label. Use other equipment only as specified herein.

CULTIVATION AFTER PLANTING

Soil may be shallow cultivated without reducing the weed control activity of Trilin. Poor weed control may result if cultivation is deeper than the treated soil since this may bring untreated soil to the surface.

CROP RECOMMENDATIONS

All recommendations are given as the broadcast rates of Trilin per acre. For band applications, decrease the amount of Trilin in proportion to the amount of surface treated per acre. Apply Trilin any time after January 1 when the soil can be worked and is suitable for good incorporation. For fall application see specific crop recommendations or "FALL APPLICATION" heading where specific crop recommendations are not given. Use the lower rate for coarser soils or soils with lower organic matter. For soils containing 10% or more organic matter do not use Trilin.

MIXING AND APPLICATION DIRECTIONS

Trilin In Water

Thoroughly clean sprayer prior to use. Fill the sprayer $\frac{1}{3}$ to $\frac{1}{2}$ full with clean water and start agitation. Add proper amount of Trilin providing sufficient agitation and finish filling the tank.

Trilin Tank Mixes in Water

For all tank mixes, continuous agitation is required (Sparger pipe agitators generally provide the best agitation in spray tanks). To prevent foaming avoid stirring or splashing air into the mixture during filling by placing the end of the fill pipe below the surface of the water in the spray tank. Do not allow the mixture to siphon back into the water source.

Mixing Order

Fill the tank $\frac{1}{3}$ to $\frac{1}{2}$ full with clean water and start agitation. Add wettable powders and dry flowables and agitate until completely dispersed. Then add aqueous suspensions (flowables): then add Trilin. Lastly, add soluble ingredients and finish filling the tank. Provide continuous agitation during filling and application. If spraying and agitation must be stopped before the tank is empty the materials may settle to the bottom. If this happens, before continuing the spray application, resuspend all of the material in the bottom of the tank. A sparge agitator is particularly useful for this purpose. It may be more difficult to resuspend the settled material than it is to suspend it originally.

Read and carefully follow all label instructions for each material added to the tank. To help assure good dispersion in the tank water, make a slurry mixture by premixing dry and flowable formulations with water and pour the slurry through a 20 or 35 mesh wet screen in the top of the tank. No finer than 50 mesh line screens in the tank should be used.

If material builds up on the walls of the spray tank, wash the tank with soapy water between fillings. Rinse and continue the spraying operation. After completion of spraying, thoroughly clean the tank lines and screens. The importance of accurate calibration and uniform application increases as the spray volume decreases. Check sprayer daily to insure proper calibration and uniform application. To insure uniform application do not apply Trilin when the wind can cause drifting of spray particles. Poor weed control may result if Trilin is applied to soils which are wet or are subject to prolonged periods of flooding.

Ground Application

Using a low pressure herbicide sprayer which will apply the spray uniformly, apply Trilin in 5 to 40 gallons of water or liquid fertilizer per acre (broadcast spray).

Aerial Application

For aerial spraying apply Trilin in 5 to 10 gallons of water or liquid fertilizer per acre. Pump pressure, nozzle arrangements, speed and height should be adjusted to provide a uniform application to the soil surface. To assure proper application spray widths use swath markers or flagmen.

Fall Application

(Areas receiving more than 20 inches average annual rainfall)

See specific crop for recommendations. Use the rates listed for spring applications for all crops for which there are no specific fall application instructions and for which Trilin is recommended as a preemergence application. Trilin should not be applied in the fall for sugarbeets, potatoes and direct-seeded tomatoes. Apply and incorporate Trilin any time between October 15 and December 31. Leave ground flat or bedded over winter. On bedded ground, beds should be knocked down to desired height prior to planting, moving some treated soil from beds into furrows. Where soil is left over winter, care should be taken not to turn up untreated soil during spring bedding operations. During seedbed preparation destroy established weeds. Destroy weeds which have become established in furrows due to uncovering of untreated soil during bedding. Trilin should not be applied in the fall to soils which are wet, are subject to prolonged periods of flooding or where rice was grown the previous year.

Plantback Restrictions

In Arizona, Colorado, Idaho, Montana, Nevada, New Mexico, Oregon, Utah, Washington and Wyoming after a spring application of Trilin, sugarbeets, red beets or spinach should not be planted for 12 months or for 14 months after fall application. Soil should be plowed to a depth of 12 inches prior to planting sugarbeets to prevent the possibility of crop injury. After a spring application, sorghum (milo), proso millet, corn or oats should not be planted for 14 months after application or for 16 months after a fall application of Trilin to avoid crop injury. If land has not been irrigated do not plant any of these crops for 18 months after a spring application or 20 months after a fall application of Trilin.

In those areas of Kansas, Nebraska, North Dakota, Oklahoma, South Dakota and Texas where at least 20 inches of rain and/or irrigation (total) was used to produce the crop, sorghum or oats should not be planted for 12 months after an application of Trilin.

Do not plant sorghum, proso millet or oats for 18 months after an application of Trilin if less than 20 inches of total water was used to produce the crop. Cool wet weather conditions during the early stage of growth may increase the possibility of injury to sorghum.

In all other areas receiving greater than 20 inches rainfall per year, moldboard plow before planting sugarbeets where a spring application of Trilin was made the previous season. Also note planting restrictions listed in the section on control of rhizome johnsongrass and other high rate programs.

For vegetables other than those listed on this label, crops should not be planted within 5 months following the application of Trilin.

PRECAUTION

Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Under normal growing conditions and if applied according to directions, Trilin will not harm the treated crop. Crop injury or soil residue may result from over application. Erratic weed control or crop injury may result from uneven application or improper soil incorporation of Trilin. Seedling disease, cold weather, deep planting, excessive moisture, high salt concentration or drought may weaken crop seedlings and increase the possibility of damage from Trilin. Under these conditions, delayed crop development or reduced yields may result.

CHEMIGATION

General Instructions

Apply Trilin only through one or more of the following types of systems: sprinkler including center pivot, lateral move, end tow, side (wheel) roll, traveler, big gun, solid set, hand move or drip irrigation systems. Do not apply Trilin through any other type of irrigation system.

Crop injury, lack of effectiveness or illegal pesticide residues in the crop may result from non-uniform distribution of treated water.

If you have questions about calibration, you should contact State Extension Service Specialists, equipment manufacturers, or other experts.

Do not connect an irrigation system used for pesticide application to a public water system unless the pesticide label prescribed safety device for public water systems is in place.

A person knowledgeable of the chemigation system and responsible for its operation, or under supervision of the responsible person, shall shut the system down and make necessary adjustments should the need arise.

Specific Instructions for Public Water Systems

Public water system means a system for the provision to the public of piped water for human consumption if such system has at least 15 service connections or regularly serves an average of at least 25 individuals daily at least 60 days out of the year.

Chemigation systems connected to public water systems must contain a functional, reduced-pressure zone, backflow preventer, (RPZ) or the functional equivalent in the water supply line upstream from the point of pesticide introduction. As an option to the RPZ, the water from the public water system should be discharged into a reservoir tank prior to pesticide introduction. There shall be a complete physical break (air gap) between the outlet end of the fill pipe and the top or overflow rim of the reservoir tank of at least twice the inside diameter of the fill pipe.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of fluid back toward the injection pump.

The pesticide injection pipeline must contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops, or in cases where there is no water pump, when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Specific Instructions for Sprinkler Irrigation Systems

The system must contain a functional check valve, vacuum relief valve, and low pressure drain appropriately located on the irrigation pipeline to prevent water source contamination from backflow.

The pesticide injection pipeline must contain a functional, automatic, quick-closing check valve to prevent the flow of liquid back toward the injection pump.

The pesticide injection pipeline must also contain a functional, normally closed, solenoid-operated valve located on the intake side of the injection pump and connected to the system interlock to prevent fluid from being withdrawn from the supply tank when the irrigation system is either automatically or manually shut down.

The system must contain functional interlocking controls to automatically shut off the pesticide injection pump when the water pump motor stops.

The irrigation line or water pump must include a functional pressure switch which will stop the water pump when the water pressure decreases to the point where pesticide distribution is adversely affected.

Systems must use a metering pump, such as a positive displacement injection pump (e.g. diaphragm pump) effectively designed and constructed of materials that are compatible with pesticides and capable of being fitted with a system interlock.

Do not apply when wind speed favors drift beyond the area intended for treatment.

Good agitation is required in the injection tank.

In moving systems, apply specified dosage of Trilin as a continuous injection. In non-moving systems inject Trilin for 15 to 20 minutes at end of cycle. Use the least amount of water possible for uniform coverage.

Mix the amount of Trilin needed for acreage to be treated into the quantity of water determined during prior calibration (refer to Mixing and Application Directions). For moving systems inject into the system continuously for one complete revolution of the field. For non-moving systems inject into system for the time established during calibration.

Stop injection equipment after treatment is completed and continue to operate irrigation equipment until all Trilin is flushed from system.

ALFALFA

Preplant

Trilin may be applied as a preplant incorporated treatment for preemergence control of labeled weeds in direct seeded alfalfa. Apply and incorporate prior to planting according to product label directions.

Broadcast Rates Per Acre

Soil Texture	Trilin (pt./Acre)
Coarse	1.0
Medium	1.0-1.5*
Fine	1.5

*Use lower range in areas receiving less than 20 inches of rainfall and irrigation.

PRECAUTION

Some crop stand reduction and stunting may occur with use of Trilin, however, reduced weed competition will allow establishment of a quality stand.

Mechanically Incorporated

Use a broadcast rate of 1.5 pints per acre on coarse soils and 2 pints on medium and fine soils (in areas receiving less than 20 inches average annual rainfall). Damage to the established alfalfa may be caused if the proper incorporation equipment is not used for thorough soil mixing. Recommended soil preparation application and incorporation instructions should be followed.

Chemigation or Water Incorporated

Trilin may be surface applied for annual grass control in established alfalfa by chemigation, or ground or aerial broadcast application equipment. Refer to "Chemigation" section of this label for use directions for chemigation.

Surface Application Activated by Rainfall or Irrigation

Broadcast surface applications of Trilin to established alfalfa may be activated by rainfall, sprinkler, flood for furrow irrigation. Rainfall or a single overhead sprinkler irrigation of 0.5 acre inch or more is required to activate Trilin. If activated by furrow irrigation, care should be taken to thoroughly wet beds between furrows. If rainfall or irrigation has not occurred within 3 days after application, Trilin may be mechanically incorporated. If mechanically incorporated, use equipment that will insure thorough soil mixing with minimum damage to the established alfalfa.

Application Timing

Applications to established alfalfa for annual grass control can be made during dormancy or semi-dormancy, or during the growing season immediately after a cutting. Because Trilin does not control established weeds, application must be made prior to the expected time of weed germination. Bromegrass and cheat begin to germinate in the fall with the onset of cooler weather. To control these weeds, apply Trilin immediately after a cutting between August 1 and October 1, but prior to weed germination. When fall applied, Trilin controls bromegrass and cheat in addition to other labeled weeds that germinate after application.

Apply Trilin at a broadcast rate of 4 pints per acre on all soil textures by chemigation or surface applied and incorporated by rainfall or irrigation.

PRECAUTION

Do not cut or graze alfalfa within 21 days after application of Trilin. Apply no more than 4 pints of Trilin during any growing season. In the growing season following application of 4 pints of Trilin to alfalfa, plant only those crops for which Trilin is registered as a preplant treatment or crop injury may occur.

ASPARAGUS Established

Follow recommended soil preparation, application and incorporation procedures for Trilin. Trilin can be applied to established asparagus as a single or as a split application. In the winter or early spring, apply to asparagus after ferns are removed but before spear emergence or apply after harvest in the late spring or early summer before turning begins. Trilin will suppress volunteer seedling asparagus and field bindweed used at the following recommended rates and application schedules.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Split Application OR	1	1.5	2
Single Application	2	3	4

In any single calendar year, the maximum Trilin to apply is 2 pints per acre on coarse soils, 3 pints on medium soils and 4 pints on fine soils.

BEAN Dry

Apply Trilin before planting using the following rates:

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. For soils with 5-10% organic matter use 2 pints on all soils.

TRILIN WITH EPTAM TANK MIX

A tank mix of Trilin and Eptam will effectively control all the following weeds in addition to those weeds listed for Trilin.

Henbit (Spring Applications)	Oat Wild
Nightshade, Black	Ragweed, Common
Nightshade, Hairy	Smartweed, Pennsylvania
Nutsedge	Velvetleaf (Buttonweed)

Follow recommended soil preparation and incorporation procedures for Trilin. Apply this tank mix from 2 days prior to planting up to planting. Incorporate immediately after application.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

PLUS

Eptam 7E* (pts.) 2.5-3.5 2.5-3.5 2.5-3.5

*To control annual grasses use Eptam 7E at a rate of 2.5 pints per acre. To control nutsedge and additional broadleaf weeds use 3.5 pints.

On soils with 2-5% organic matter use Trilin at 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. For soils with 5-10% organic matter use 2 pints of Trilin on all soils.

PRECAUTION

Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Do not use this tank mix on some blackeyed peas (beans), lima beans, soybeans, mung beans and other flatpodded beans except Romano. Do not use foliage from treated plants for feed, forage or for grazing.

Fall application in Idaho, Oregon and Washington

Apply and incorporate Trilin between October 15 and December 31. Use a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils and 1.5 pints on fine soils. Destroy established weeds during seedbed preparation.

BEAN Guar and Mungbean

Apply and incorporate Trilin prior to planting at 1 pint per acre on coarse soils and 1.5 pints on medium and fine soils.

BEAN Lentil (Except CA)

Apply and incorporate Trilin before planting. Plant no deeper than 1.5 inches when temperature and moisture in seedbed promote fast germination and emergence.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Spring Application	1	1	1.5
Fall Application	1.25	1.25	1.75

For soils with 2-5% organic matter use 1.5 pints per acre.

PRECAUTION

Lentil tolerance to trifluralin is marginal. Injury may occur under conditions of plant stress such as cold weather, low fertility and disease or insect damage. Spring application may increase the probability of stand reductions.

BEAN Snap and Lima

Apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils.

CARROT

Apply and incorporate Trilin prior to planting.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

CASTOR BEAN

Apply and incorporate Trilin before or immediately after planting. If applied and incorporated after planting, set equipment so as to not disturb the seed.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

CELERY

Apply Trilin as a soil incorporated treatment. Trilin may be applied to direct-seeded or transplant celery before planting, at planting or immediately after planting.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.25-1.5	1.5-2.0

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

Use the lower rate in the range in areas receiving less than 20 inches total annual rainfall and/or irrigation.

CHICORY

Trilin may be applied as a preplant incorporated treatment to chicory grown either as a root crop or leafy vegetable

Cichorium intybus

Considered to be a root crop, may yield the following:

Chicory - the dried and processed root used as a coffee substitute.

Radichio - green leaves harvested from field grown plantings.

Belgian Endive - white leaves grown in the dark growth from field grown rootstalks.

Cichorium endiva

Considered to be a leafy vegetable, may yield the following:

Escarole - curly green leaves from field grown plantings.

Endive - very curly green leaves from field grown planting.

Apply Trilin as a soil incorporated treatment in spring or early summer prior to planting,

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium soils and 2.0 pints on fine soils. On soils with 5-10% organic matter use 2.0 pints on all soil textures.

COLE CROPS

Broccoli, Brussels Sprout, Cabbage & Cauliflower

Transplant

Apply and incorporate Trilin prior to transplanting only.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

Direct-Seeded

Use Trilin before planting at a broadcast rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils and soils with 2-5% organic matter. Direct-seeded cole crops have exhibited marginal tolerance to recommended rates of Trilin.

Warning: Stunting or reduced stands may occur.

COTTON

Apply Trilin before planting, immediately after planting, or at lay-by.

Preemergence

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

Use 1.5 pints per acre on coarse and medium textured soils, 2 pints on fine soils with 2-5% organic matter and 2 to 2.5 pints on all soils with 5-10% organic matter.

Postplant

Do not disturb the seed when incorporating Trilin postplant.

Lay-by

Trilin can be applied and incorporated any time up to lay-by, but not less than 90 days before harvest. Direct lay-by applications onto the soil between the rows and beneath emerged cotton plants at the same rates as for a preemergence application.

Fall Application

Trilin may be applied and incorporated any time between October 15 and December 31. The ground may be left flat or bedded-up over winter. Where soil is left flat, take care not to turn up untreated soil from beds into furrows. On bedded ground, knock down beds to desired heights before planting, moving some treated soil into the furrow from the beds. Destroy established weeds during preparation of seedbed. Before planting, destroy weeds which may have become established in furrows during the uncovering of untreated soil during bedding. Trilin should not be applied to wet soil or soils which are subject to prolonged periods of flooding.

Application Over Standing or Shredded Cotton Stalks

Trilin may be broadcast applied, after September 1, over the top of standing cotton stalks or after shredding using ground or aerial equipment. Trilin must be incorporated once within 24 hours after application using a disc operated at 4 to 6 mph. The second incorporation may be delayed until spring prior to bedding. To avoid dilution of the herbicide or bringing untreated soil to the surface, the treatment should not be chiseled, ripped or deep plowed following incorporation.

Broadcast Rates Per Acre – Fall Application Only

In Alabama, Arkansas, Northern Florida, Georgia, Louisiana, Mississippi, Southeastern Missouri, North Carolina, New Mexico, Oklahoma, South Carolina, Tennessee and Texas, apply and incorporate Trilin at a broadcast rate of 2 pints per acre on coarse and medium soils and 2.5 pints on fine soils.

In Arizona, California and Nevada, a broadcast rate of 1.5 pints per acre should be used on coarse soil, 2 pints on medium soil and 2.5 pints on fine soil.

For other states where cotton may be grown, apply Trilin at a broadcast rate of 1 pint per acre on coarse soils, 1.5 pints on medium soils and 2 pints on fine soils. For coarse soils with 2-5% organic matter use 1.5 pints. For soils with 5-10% organic matter use 2 to 2.5 pints.

Incorporation of Trilin With Bedding Implements for Weed Control in Cotton

Bedding implements (listers and hippers) may be used to soil incorporate Trilin for weed control in cotton. Because bedding implements do not provide thorough soil mixing under all conditions, it is important to closely follow use directions to optimize weed control. Weed control resulting from single pass incorporation with bedding equipment will be reduced compared to conventional double pass incorporation. Use the application rate recommended for the soil texture to be treated.

Soil Preparation

Crop Residues or Existing Weeds: Ground cover, such as crop residues or existing weeds, can interfere with uniform soil incorporation of Trilin. A manageable level of such ground cover will allow uniform incorporation into the top 2 to 3 inches of soil. Ground cover or crop residues, if excessive, should be reduced by appropriate soil tillage prior to the application of Trilin.

General Soil Conditions

The soil surface should be smooth enough to allow for uniform application and efficient incorporation of Trilin. Apply Trilin when soil moisture is sufficient to allow the breakup of large clods and uniform mixing during the incorporation process.

Use Directions for Bedding Equipment

A lister or disc bedder may be used to incorporate Trilin. Operate the implement according to the manufacturers use directions in order to produce beds of the desired height. A ripper shank, sweep or chisel shank should be mounted on the bedder in a position behind the spray nozzles but ahead of the bedder tool to help distribute Trilin in the center of the bed. The use of bed tillage equipment such as rolling cultivators, P.T.O. driven rod weeders or bed conditioners after the bedding operation will provide additional soil mixing. Avoid deep tillage which might bring untreated soil to the surface resulting in loss of weed control. Weather conditions, cultural practices, bed tillage and planting procedures can affect the distribution of Trilin treated soil. Weed control obtained will be dependent upon how uniformly Trilin treated soil is distributed over the soil surface at the time of planting.

If Trilin treated soil is moved, exposing untreated soil, during bed tillage or planting, a band application of Trilin at planting or a postemergence application may be required to restore uniform weed control.

PRECAUTION

Do not incorporate with bedding equipment if the soil is too wet for uniform soil mixing.

Trilin for Weed Control in Conservation Tillage Cotton

Trilin may be applied and incorporated before planting, after planting prior to crop emergence, or at lay-by for weed control in conservation tillage cotton. Single or multiple application may be made so long as maximum application rates are not exceeded and rotational crop restrictions are followed.

Strip Planting into Small Grain Cover Crops

Fall planted cover crops may be utilized to control wind erosion and protect developing crop seedlings from wind damage. Prior to planting cotton, the cover crop may be treated with a contact herbicide to prevent continued growth and development and prevent competition with crop seedlings for water and soil nutrients. The standing cover crop (now dead) continues to control wind erosion and provide protection to the developing crop until it is well established.

In strip planting, cotton is seeded into competition free bands established in the cover crop. Competition-free bands may be established by leaving unseeded drill rows when seeding the cover crop, or by tillage of use of a contact herbicide to prepare competition-free bands prior to planting.

Application Before or After Planting (Within Competition-Free Bands)

Apply Trilin using low pressure ground equipment as a band (within the weed free zone) or as a broadcast treatment. Application and incorporation may occur before planting or after planting prior to crop emergence. If applied after planting, set incorporation equipment so as to not disturb the planted seed.

Incorporation

Equipment should be adapted to the width of the competition-free band. Use equipment that will uniformly mix Trilin into the weed germination zone. Weed control resulting from single pass incorporation or with incorporation equipment that does not result in thorough mixing of Trilin treated soil may be reduced compared to conventional double pass incorporation. Implements used to incorporate Trilin after planting should be operated so that they do not disturb the planted seed or growing crop.

Broadcast Rates Per Acre

Use 1.0 to 2.0 pints per acre on coarse, 1.5 to 2 pints on medium and 2 to 4 pints on fine soils. Use the lower rate in the rate range when additional sequential applications of Trilin are anticipated. Use the higher rate in the rate range where high crop residues are present, and where dense weed populations are anticipated.

For band treatments, reduce the application rate in proportion to the row spacing and band width treated. For example, treating a 12-inch band where the row spacing is 36 inches would require $\frac{1}{3}$ of the recommended broadcast rate per acre (12 inches divided by 36 inches = $\frac{1}{3}$).

Lay-by Applications

Lay-by applications may be made in established cotton from the four true leaf stage of growth up to lay-by, but not less than 90 days before harvest. Apply Trilin uniformly to the soil surface using drop nozzles if necessary. Apply at a broadcast rate of 1 pint per acre for coarse, 1.5 pints per acre for medium and 2 pints per acre for fine textured soils. Soil incorporate using one pass of a sweep-type cultivator or properly adjusted rolling cultivator. Operate cultivation equipment at speeds sufficient to provide vigorous soil mixing and exercise care to avoid mechanical injury to the crop. Cumulative lay-by application rate may not exceed the lay-by application rate given for each soil texture.

Chemigation

Trilin may be applied through properly equipped chemigation systems for weed control in conservation tillage cotton. Apply the recommended rate of Trilin in 0.5 to 1.0 acre in the sprinkler irrigation. Apply Trilin before planting or after planting, but prior to weed or crop germination. Trilin does not control established weeds. Soil incorporation is not required when applied by chemigation. Soil treated with Trilin may be shallow-cultivated without reducing weed control activity. See "Chemigation" section for detailed use directions. Apply at a broadcast rate of 1 to 3 pints per acre for coarse, 1.5 to 4 pints per acre for medium and 2 to 4 pints per acre for fine textured soils. Use the lower rate in the rate range when additional sequential applications of Trilin are anticipated. Use the higher rate in the rate range when high crop residue levels are present, where dense weed populations

are anticipated, or where no additional subsequential applications of Trilin are to be made (See Sequential Applications section).

Sequential Applications

Trilin may be applied one or more times sequentially during the growing season using the rates and methods of application described for full season weed control. The maximum dosage that can be used for a single application cannot exceed the rates shown for each application method. The maximum cumulative application rate that may be applied within the same growing season cannot exceed 4 pints per acre.

Crop Rotation

See the Plantback Restrictions section for specific rotational crop restrictions. When the cumulative application rate exceeds the application rate of 1.5 pints per acre for coarse, 1.5 pints per acre for medium and 2 pints per acre for fine textured soils, plant only those crops for which Trilin can be applied as a preplant incorporated treatment in the season following the Trilin application or crop injury may result.

Small grain cover crops that will not be grazed or harvested and are intended for prevention of wind erosion in conservation tillage cotton may be planted in the fall following spring applications of up to 4 pints per acre Trilin. Injury in the form of reduced stands or delayed emergence and development may result when small grains are planted under these conditions.

SPECIAL USE DIRECTIONS FOR COTTON

Fall Panicum

Apply and incorporate Trilin broadcast at the rate of 2 pints per acre on coarse and medium soils.

Texas Panicum

Apply and incorporate at the labeled use rate for the soil type. Uniformly incorporate to a depth of at least 3 inches. Adequate soil moisture is essential for herbicide activation. Any deviation from these recommendations can result in reduced control.

Rhizome Johnsongrass

Rhizome johnsongrass can be controlled in all cotton producing states except Arizona by using a double rate program applied for 2 consecutive years as follows.

Soil Preparation

For satisfactory results proper soil preparation is essential. To bring rhizomes to the top of the soil use a chisel plow or similar implement. Follow with a disc twice before application to cut rhizomes into small 2 to 3 inch pieces. Any emerged johnsongrass should be destroyed.

Application

Choose one of the following programs which best meets your cultural practices:

For spring application apply Trilin prior to planting in the spring for 2 consecutive years. A broadcast rate of 2 pints per acre should be used on coarse soils, 3 pints on medium soils and 4 pints on fine soils.

OR

For fall application use Trilin between October 15 and December 31 for 2 consecutive years at the same rates as a spring application.

Incorporation

For good rhizome johnsongrass control deep incorporation is necessary. Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two passes are necessary with the second pass in a different direction from the first.

Cultivation

To remove johnsongrass plants which have escaped control, timely cultivations during the crop season are necessary to obtain effective control. Effective control cannot be obtained with only 1 year of double rate Trilin use.

Crop Rotation

Plant only rice and those crops for which Trilin can be applied as a preplant treatment following a double rate treatment season or injury may result.

Pigweed and Seedling Johnsongrass Control

In Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Southeastern Missouri, North Carolina, South Carolina, Tennessee and Southern Virginia, apply Trilin preplant at a broadcast rate of 1 to 1.5 pints per acre on coarse soils, 1.5 to 2 pints on medium soils and 2 pints on fine soils. **Exception:** Use 3 pints per acre on fine soils in Louisiana.

Additional Weed and Grass Control

In the Texas Gulf Coast counties of Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton, apply Trilin at a broadcast rate of 1.5 pints per acre on coarse soils, 2 pints on medium soils and 3 pints on fine soils 2 weeks prior to planting.

PRECAUTION

When using higher usage rates, plant cotton after early season adverse weather conditions to avoid additional stress to the cotton plants due to cool, wet weather early in the growth cycle, which could cause reduced stands, delayed maturity and reduced yields.

TANK MIXES AND OVERLAYS

Observe all precautions and limitations on the label of each product used in tank mixes and overlays.

TRILIN WITH COTTON-PRO – Arizona, New Mexico and West Texas TRILIN WITH CAPAROL 4L – Arizona, New Mexico and the upper and lower El Paso Valley, Texas

Trilin with Cotton-Pro or Caparol 4L will control certain grasses and broadleaf weeds listed for Trilin alone plus the following weeds.

Annual Morningglory	Prickly Sida (Teaweed)
Groundcherry (Annual)	Ragweed
Malva	Smartweed
Mustard	Wild Oat

Shallow germinating seedlings of cocklebur and coffeeweed will also be controlled.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.25-1.5	2
PLUS			
Cotton-Pro (pts.)	3.125	4	4
OR			
Caparol 4L (pts.)	2.4-3.2*	4	4

Do not use on sands and loamy sands. Use proportionally less for band application. *Use less than 3.2 pints per acre only in Arizona.

Mixing Directions

Add Cotton-Pro or Caparol 4L to a partially filled tank of water. Add Trilin and fill tank. During the filling and spraying operation, agitate continuously. If bypass agitation is used, minimize foaming by hanging the bypass line stop at the bottom of the tank. Avoid leaving the spray mixture in the tank without constant agitation.

Crop Rotations

After a spring application of Trilin plus Cotton-Pro or Caparol 4L, cabbage, okra, onions and peas may be planted in the fall. Winter barley, winter rye and winter wheat, if plowed down and not used for food or feed, can be planted in the fall also. Refer to the Cotton-Pro or Caparol 4L label for directions, cautions and precautions.

PRECAUTION

Do not use Trilin with Cotton-Pro or Caparol 4L tank mix in the cut areas of newly leveled fields, in areas of excess salt or where flooding over the beds is likely to happen. Do not plant cotton in tractor wheel depressions. These conditions may cause crop injury. On mulch planted cotton, water back only after cotton seedlings are well-established. Do not feed foliage from treated plants to livestock or graze treated areas.

TRILIN WITH METURON 4L, METURON 80DF, COTORAN 4L OR COTORAN DF (Except in Arizona)

Follow recommended soil preparation and incorporation procedures for Trilin. A tank mix of Trilin with Meturon 4L, Meturon 80DF, Cotoran 4L or Cotoran DF effectively controls all the annual grasses and broadleaf weeds listed for Trilin alone plus these additional weeds:

Buttonweed	Morningglory	Sesbania
Cocklebur	Prickly Sida (Teaweed)	Sicklepod
Groundcherry, Wright	Ragweed	Smartweed
Jimsonweed	Ryegrass	Tumbleweed

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2
PLUS			
Meturon 4L (pts.)	2	3.125	4
OR			
Meturon 80DF (lbs.)	1.25	1.9	2.5
OR			
Cotoran 4L (lbs.)	2	3.125	4
OR			
Cotoran DF (lbs.)	1.2	2	2.4

Mixing Directions

Add Meturon or Cotoran to a partially filled tank of water. Add the Trilin after Meturon or Cotoran is thoroughly mixed and continue filling. Agitate continuously throughout the filling and application operations. Do not leave spray mixture in tank without constant agitation. If bypass agitation is used, minimize foaming by hanging the bypass line stop at the bottom of the tank. Apply in 15 to 40 gallons of water per acre.

West Texas Only

Do not use tank mix of Trilin plus Meturon or Cotoran on sandy, loamy sand or fine sandy loam soils. Do not use on cotton planted in furrows.

Arkansas, Louisiana and Mississippi Only

On sandy loam soils low in organic matter use 1.5 pints Meturon 4L or Cotoran 4L or 1 pound Meturon 80DF or Cotoran DF in tank mix with Trilin.

New Mexico

Do not plant treated land with crops other than cotton until 1 year after the last application. Do not use on sandy or coarse textured soils of less than 1% organic matter.

PRECAUTION

Crop injury may result if treated land is planted to anything but cotton within 6 months of the application of Trilin plus Meturon or Cotoran. Do not feed foliage from treated cotton plants or gin trash to livestock. Do not mix Trilin plus Meturon or Cotoran with liquid fertilizer.

TRILIN PREPLANT FOLLOWED BY METURON 4L, METURON 80DF, COTORAN 4L OR COTORAN DF OVERLAY

Apply and incorporate Trilin as recommended for the specific soil texture. Apply Meturon 4L or Cotoran 4L at 2 to 4 pints per acre or Meturon 80DF or Cotoran DF at 1.2 to 2.4 pounds per acre as a preemergence surface treatment. Use the lower rate on light silt and sandy soil low in organic matter.

PRECAUTION

Observe all precautions and limitations on the labels of each product used in tank mixes and overlays.

TRILIN PREPLANT FOLLOWED BY DIREX 4L OR KARMEX DF OVERLAY (East of the Mississippi River plus Arkansas, Southeastern Missouri, Louisiana and Eastern Texas)

Incorporate Trilin prior to planting. Follow with a preemergence application of Direx 4L or Karmex DF. This will effectively control all the weeds controlled by Trilin alone plus these additional weeds.

Annual Ground, Cherry	Pennycress	Velvetgrass
Annual Morningglory	Ragweed	Wild Lettuce
Dogfennel	Shepherdspurse	Wild Mustard

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2
PLUS			
Direx 4L (pts.)	0.5	1	1.75
OR			
Karmex DF (lb.)	0.33	0.67	1

PRECAUTION

Observe all precautions and limitations on the label of each product used in tank mixes and overlays. Direx 4L or Karmex DF should not be used on soils with less than 1% organic matter as crop injury may result. Do not feed foliage from treated plants to livestock or graze treated areas.

TREES GROWN FOR PULPWOOD

Aspen, Cottonwood and Poplar

New Plantings

Apply and incorporate Trilin before planting.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.25-1.5	1.5-2

For soils with 2-5% organic matter use 1.5 to 2 pints. On soils with 5-10% organic matter use 2 pints. Use lower rate in areas receiving less than 20 inches total rainfall and irrigation.

Established Plantings

Apply as a directed spray to the soil and use incorporation methods not injurious to the crop.

Broadcast Rates Per Acre

	All Soil
Trilin (pts.)	2-4

Application rate within the rate range may be adjusted according to weed pressure.

Johnsongrass Suppression

Proper soil preparation before application is necessary for satisfactory results. Use a chisel plow or similar implement to bring rhizomes to the soil surface. Then work the soil twice using a tandem disc to cut rhizomes into small (2 to 3 inch) pieces and to destroy emerging johnsongrass.

Broadcast Rates Per Acre

	All Soil
Trilin (pts.)	4

Incorporation: Incorporate twice with a tandem disc set to cut 4 to 6 inches deep and operated at 4 to 6 mph.

Cultivation: Some johnsongrass plants will escape. Timely cultivation with tillage implements or spot spraying with effective postemergence herbicides will improve the level of johnsongrass control.

CUCURBITS

Cantaloupe, Cucumber and Watermelon

Postplant Emerged in Western United States including Texas

Apply Trilin as a directed spray to the soil between the rows and beneath plants which are in the three to four true leaf stage.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

EGGPLANT

(Except CA)

Apply and incorporate Trilin before transplanting. Incorporate to a depth of 3 inches. Do not make more than one application per season.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints.

PRECAUTION

Avoid transplanting until soil temperatures have warmed in late spring.

FIELD CORN, GRAIN SORGHUM AND CORN FODDER, FORAGE AND SILAGE

Apply Trilin to field corn or grain sorghum (8 inches or taller) as an over-the-top or directed spray to effectively control weeds listed for Trilin. Trilin applied as an over-the-top spray or as a directed spray in field corn and grain sorghum will control shattercane in addition to those other weeds listed on the label for Trilin.

Soil Preparation

Cultivate before a Trilin application to insure loose, friable soil, to remove established weeds and to cover the base of plants with soil.

Application Directions

Trilin should be applied and incorporated at the recommended rates for the soil texture when the crop is well established (8 inches or taller). Trilin may be applied either as an over-the-top spray or as a directed spray. Drop nozzles should be used if foliage prevents uniform coverage of soil surface. Soil incorporation may be accomplished with only one pass of a sweep-type cultivator or a properly adjusted rolling cultivator. The sweep-type cultivator should have three to five sweeps per row middle and be operated at 6 to 8 mph. Set the middle sweeps so as to avoid exposing untreated soil. Adjust the incorporation tools to prevent crop injury.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	0.75-1	1-1.5	1.5-2

Corn Only

Apply 1 to 1.5 pints per acre in Alabama, Florida, Georgia, North Carolina, South Carolina and Virginia to control fall panicum and Texas panicum.

TRILIN WITH ATRAZINE TANK MIX

Trilin can be tank mixed with atrazine for additional weed control in field corn and grain sorghum.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	0.75-1	1-1.5	1.5-2
PLUS			
AAtrex 4L (pts.)	2.4	4.75	6

When using AAtrex 4L use the rates listed above. For other atrazine formulations, use equivalent rates. When using AAtrex NineO 1 pint of AAtrex 4L equals 0.55 pound of NineO. One pint of AAtrex 4L equals 0.62 pound of Atrazine 80W.

Do not use the above tank mix on coarse soils for grain sorghum.

Apply and incorporate the Trilin/Atrazine tank mix as directed on the Trilin label for field corn and grain sorghum.

PRECAUTION

Do not apply Trilin to sweet corn or corn grown for seed. Do not apply to corn or sorghum as a preplant or preemergence treatment or crop injury may occur. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays.

FLAX

Fall Application

Trilin may be applied and incorporated in the fall for weed control in spring seeded flax. Ground cover from existing weeds or previous crop should be at a manageable level so that there is no interference with incorporation.

Apply at the broadcast rates per acre of 1 pint on coarse soil, 1.5 pints on medium soil and 2 pints for fine soils.

Incorporation Directions

Incorporate one time within 24 hours after application. The second incorporation should be performed in the spring prior to seeding flax. The incorporation operations should result in a thorough mixing of Trilin with soil. Otherwise, erratic weed control may result.

Incorporation Equipment

Follow recommended soil incorporation procedures for Trilin. Incorporation operations or any other tillages performed in the spring prior to seeding should be relatively shallow so as to maintain a firm seedbed, and the seedbed should be packaged just prior to seeding. Seeding should be done with a press drill or hoe drill. Seed into a moist seedbed no more than 1.5 inches deep. Do not delay the first incorporation more than 24 hours after application.

FRUIT AND NUT CROPS AND VINEYARDS

For areas receiving greater than 20 inches average annual rainfall

On new plantings of citrus, pecan trees and vineyards apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse soils, 1.5 pints on medium soils and 2 pints on fine soils. On soils with 2-5% organic matter use 1.5 pints on fine soils and on soils with 5-10% organic matter use 2 pints. Use Trilin at a broadcast rate of 2 to 4 pints per acre for all soil textures for non-bearing established plantings of citrus and pecan trees and bearing plantings of grapefruit, lemon, orange, pecan, tangelo and tangerine trees.

For areas receiving less than 20 inches average annual rainfall

On new plantings of almond, apricot, citrus, nectarine, peach, pecan and walnut trees apply and incorporate Trilin prior to planting at a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils, 1.5 pints on fine soils, on soils with 2-5% organic matter use 1.5 to 2 pints and 2 pints on soils with 5-10% organic matter.

New Plantings of Vineyards

Apply and incorporate Trilin prior to planting at a broadcast rate of 1 to 1.5 pints per acre on coarse soils, 1.5 to 3 pints on medium soils and 3 to 4 pints on fine soils or soils with 2-10% organic matter. Do not use more than 2 pints per acre on heat-treated grape rootings.

Postplant Application

For postplant applications on bearing or non-bearing established plantings of vineyards and almond, apricot, grapefruit, lemon, nectarine, orange, peach, plum, prune, tangelo, tangerine and walnut trees. Use Trilin at a broadcast rate of 2 to 4 pints per acre for all soil textures. Do not apply to vineyards within 60 days of harvest. In established plantings, use Trilin as a directed spray to the soil. Use incorporation methods not injurious to the trees or vines.

Rhizome Johnsongrass Control

For areas receiving less than 20 inches average annual rainfall. Control rhizome johnsongrass with postplant applications in bearing and non-bearing established plantings of vineyards and almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, tangelo, tangerine and walnut trees by applying Trilin for 2 consecutive years.

Soil Preparation

Soil should be worked thoroughly to bring the rhizomes near the surface.

Application

Use Trilin at a broadcast rate of 2 quarts per acre on all soil textures each year for 2 consecutive years. Do not apply to vineyards within 60 days of harvest.

Incorporation

Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary using a different direction for the second.

Cultivation

As some johnsongrass plants will escape, cultivation is necessary to obtain effective control. Effective control cannot be obtained with only 1 year of Trilin use.

PRECAUTION

Crop injury may result if the 2 quart rate is used on new plantings. Orchard or vineyards should not be interplanted with other crops. If the treated vineyards and orchards are diverted to other crop uses, plant only those crops for which Trilin has been registered as a preplant treatment for the next cropping season.

Bindweed control in California

Use Trilin for the control of field bindweed in vineyards and for almond, apricot, grapefruit, lemon, nectarine, orange, peach, pecan, tangelo, tangerine and walnut trees.

Use Trilin at a broadcast rate of 4 pints per acre on all soil textures. Trilin must be applied in the spring with a specially designed spray blade which applies a thin concentrated layer at a soil depth of 4 to 6 inches. This layer of Trilin prevents bindweed shoots from emerging.

Land Preparation

All weeds and grasses should be destroyed with soil tillage prior to applying Trilin. This tillage is necessary to prevent trash from interfering with the operation of the spray blade.

Equipment

A spray blade capable of running 4 to 6 inches below the surface of the soil should be used. The spray blade should be equipped with nozzles located

under the blade and directed so that the Trilin spray will be trapped under the soil which is flowing over the blade as it is pulled through the soil. A sufficient number of nozzles should be used with spacing that will uniformly apply the Trilin underground in a thin horizontal layer.

Application

Use Trilin in 40 to 80 gallons of water per acre. Operate the spray blade at a depth of 4 to 6 inches.

PRECAUTION

After rainfall or irrigation some soils may crack as they dry. Bindweed may emerge if the cracks extend through the Trilin layer. Prevent or eliminate cracks by shallow discing or other tillage. Avoid deep tillage which disturbs the subsurface layer. Cultivation or tillage also aids the control of germinating seeds.

GREENS

Turnip greens grown for processing, Collards, Kale and Mustard Greens

Apply and incorporate Trilin prior to planting at 1 pint per acre on coarse soils and 1.5 pints on medium and fine soils.

HOPS

Apply and incorporate Trilin while the crop is dormant using a broadcast rate of 1 pint per acre on coarse soils, 1.25 pints on medium soils and 1.5 pints on fine soils. Use incorporation equipment that will insure thorough soil mixing with minimum damage to the crop.

IRRIGATION WATER RINGS NON-BEARING CITRUS TREES (Except CA)

Apply to non-bearing citrus trees through irrigation water rings to provide preemergence weed control. Mix at a rate of 12 fluid ounces of Trilin per 500 gallons of water. Agitate until uniformly dispersed in tank.

Apply 10 gallons of the mixture per 4 foot diameter water ring per tree. Trilin should be applied at the second or third watering and should not be applied in combination with any other pesticide.

KENAF

Ground Application

Using a low pressure herbicide sprayer which will apply the spray uniformly, apply Trilin in 5 to 40 gallons of water (broadcast spray).

Aerial Application

For aerial spraying apply Trilin in 5 to 10 gallons of water. Pump pressure, nozzle arrangements, speed and height should be adjusted to provide a uniform application to the soil surface. To assure proper application spray widths use swath markers or flagmen.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	0.75-1	1-1.5	1.5-2

Incorporation Directions

General Directions

Erratic weed control and/or crop injury may result if directions are not followed. The soil surface should be free of existing weeds and excessive trash or clods.

Before Planting

Incorporation of Trilin must take place within 24 hours after application and may be accomplished in one pass with a sweep type or a rolling cultivator. Follow recommended soil incorporation procedures for Trilin.

PRECAUTION

Without reducing the weed control activity of Trilin, soil treated with Trilin may be shallowly cultivated, rotary hoed or handhoed. Do not cultivate deeper than (1 to 2 inches) the Trilin treated layer of soil as this may bring untreated soil to the surface and poor weed control may result. Kenaf treated with Trilin must not be used as a food, forage or feed.

LUPINE (Except CA)

Apply and incorporate before planting.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium soils and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints.

MUSTARD

Grown for seed or processing for food in Minnesota, Montana and North Dakota. Apply and incorporate Trilin prior to planting at 1 pint per acre on coarse soils and 1.5 pints on medium and fine soils.

NUT CROPS

See FRUIT AND NUT CROPS AND VINEYARDS for instructions.

OKRA

Apply and incorporate Trilin before or immediately after planting. If applied and incorporated after planting, set equipment so as to not disturb the seed.

Broadcast Rates Per Acre

Trilin (pts.)	Coarse	Medium	Fine
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

ONIONS

Grown for Dry Bulbs Only

Apply Trilin as a directed spray to the soil between the onion rows. Avoid spraying the onion tops or exposed bulbs. Do not apply within 60 days of harvest.

Broadcast Rates Per Acre

Trilin (pts.)	Coarse	Medium
	0.75-1	1-1.25

For band applications, use proportionately less Trilin. Use the lower rates where light weed pressure is anticipated.

Soil incorporation may be accomplished by operating sweep-type or rolling cultivators 2 to 4 inches deep at 6 to 8 mph. Incorporation equipment must mix Trilin uniformly in the soil. Incorporate with two passes. The first pass must be within 24 hours of application or erratic weed control may result. Avoid covering exposed onion bulbs with treated soil during incorporation as injury to the crop may occur. Care should be taken to avoid injury to the roots during incorporation.

PRECAUTION

Applied according to these directions and under normal growing conditions Trilin will not adversely affect onions. Diseases, improper incorporation depth, excessive moisture, high salt concentration or drought may weaken the crop and increase the possibility of damage from Trilin. Under these conditions, delayed crop development or reduced yields may result.

ORNAMENTALS

(Except CA)

Apply and mechanically incorporate Trilin prior to planting new nursery stock liners, ornamentals, trees and woody shrubs and gladioli. Gladioli corms less than 1 inch diameter may be injured by preplant applications. Trilin may also be applied to these and other listed ornamentals (see below) after they are established. When mechanically incorporated after planting, the implement should be adjusted so that treated soil is thrown toward and around the plants in the row.

Broadcast Rates Per Acre

Trilin (pts.)	Coarse	Medium	Fine
	1	1.5	2

For the indicated ornamental groundcovers, apply 1 gallon per acre (3 ounces per 1000 square feet) of Trilin in 5 to 40 gallons of water per acre and incorporate within 24 hours with at least a ½ inch rain or its equivalent in sprinkler irrigation.

Woody Shrubs	Trees	Ground Cover
Andromeda, Japanese	Almond	Aaronsbeard
Arborvitae, American	Apple, Crabapple	Bellflower, Adriatic
Azalea	Apricot	Bellflower, Poscharsky
Barberry, Japanese	Ash, White	Ceanothus
Barberry, Mentó	Bald Cypress	Coreopsis
Boxwood, Common	Birch, European White	Cotoneaster
Boxwood, Harlands	Black gum	Coyote Brush
Boxwood, Littleleaf	Cherry	Crown Vetch
Camellia, Japanese	Chestnut, Chinese	Daisy Trailing African
Camellia, Sasanqua	Cottonwood	Fern, Asparagus
Cherry Laurel, American	Dogwood, Flowering	Gazania
Cinquefoil	Dogwood, Kousa	Germander
Cleyera, Japanese	Douglas Fir	Ice Plant, Largeleaf
Cotoneaster, Cranberry	Fir, Balsam	Ivy, Algerian
Cotoneaster, Zabel	Hemlock, Canada	Ivy, English
Deutzia	Honey Locust	Lily-of-the-Nile

Woody Shrubs

Elaegmus, Silverberry
Euonymus, Spreading
Euonymus, Winged
Euonymus, Wintercreeper
Firethorn
Forsythia
Guava, Pineapple
Holly
Honeysuckle
India Hawthorn
Juniper
Laurel, Mountain
Lilac, Common
Mock Orange
Pittosporum, Japanese
Privet
Red Cedar, Eastern
Rhododendron
Spiraea, Vanhoutte
Viburnum
Weigela
Willow
Yew, Anglojap
Yew, Japanese
Yew Pine

Trees

Larch, Japanese
Locust, Black
Maple, Norway
Maple, Red
Maple, Silver
Maple, Sugar
Oak, Pin
Oak, Red
Oak, Scarlet
Peach
Pine, Austrian
Pine, Easter White
Pine, Japanese Black
Pine, Loblolly
Pine, Red
Pine, Scotch
Planetree, London
Plum
Redbud, Eastern
Spruce, Colorado
Spruce, Norway
Spruce, White
Sweet Gum
Sycamore
Tulip Tree
Walnut, Black

Ground Cover

Lilyturf, Bigblue
Marigold
Myoporum
Plumbago, Dwarf
Rockrose
Rosemary
Rupturewort
Snow-in-Summer
Speedwell
St. Johnswort
Stoncrop (Sedum)
Strawberry, Beach
Thrift
Verbena
Wirevine, Creeping
Yarrow, Wolly
Zoysiagrass

Roses and Other Established Flowers

African Daisy	Gaillardia	Rose
Aster (perennial)	Gladiolus	Salvia
Balsam	Golden Glow	Shasta Daisy
Black-eyed Susan	Impatiens	Snapdragon
Calendula	Ixora	Snow-on-the- mountain
Carnation	Lobelia	Stock
Centaurea, Velvet	Lupine	Sunflower
Chrysanthemum	Marigold	Sweet Alyssum
Coreopsis	Marigold, Cape	Sweet Pea
Cornflower	Morningglory	Sweet Sultan
Cosmos Dahlia	Nasturtium	Sweet William
Dianthus	Petunia	Vinca
Dusty Miller	Phlox	Yarrow
Floss Flower	Pincushion Flower	Zinnia
Forget-me-not	Poppy, California	
Four O'Clock	Portulaca	

UNDER PAVED SURFACES

General Use Instructions and Site Preparation:

Trilin should be used only where the area to be treated has been prepared according to good construction practices. If rhizomes, stolens, tubers or other vegetative plant parts are present in the site, they should be removed by scalping with a grader blade to a depth sufficient to insure their complete removal.

Application should be made only when final grade is established or after additions of base rock. Do not move soils following Trilin application and do not apply Trilin to areas where asphalt is to be laid directly on top of soil. Paving should follow Trilin applications as soon as possible.

Large Areas: Apply Trilin in sufficient water to insure thorough wetting of the soil surface or penetration of the spray solution through the base rock layer. A minimum of 150 gallons per acre is recommended. Apply with any sprayer that will apply the spray uniformly. Add the recommended amount of Trilin to clean water in the spray tank during the filling operation. Agitate before spraying.

Small Areas: For treating small areas, a tank type hand sprayer or sprinkling can may be used. Before application determine the amount of water and Trilin necessary to uniformly cover the area to be treated. Shake or stir the spray solution prior to application.

	Ounces Per 1000 Square Feet	Gallons Per Acre
Trilin	9-12	3-4

PEA Dry and English

Trilin Alone

Apply and incorporate Trilin prior to planting at a rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils.

Fall application in Idaho, Oregon and Washington

Apply and incorporate Trilin any time between October 15 and December 31 using a broadcast rate of 1 pint per acre on coarse soils, 1.25 to 1.5 pints on medium soils and 1.5 pints on fine soils. Destroy established weeds during seedbed preparation. Do not apply Trilin in the fall to soils which are wet or are subject to prolonged periods of flooding.

TRILIN WITH FAR-GO TANK MIX in Idaho, Oregon and Washington

Trilin plus Far-Go controls wild oat in addition to other annual grasses and broadleaf weeds controlled by Trilin.

Application Rates

Broadcast 0.75 pint of Trilin per acre on coarse and medium soils, 1 pint on fine soils. Use 1.25 quarts of Far-Go per acre for all soil textures.

Incorporation Directions

Apply and incorporate up to 3 weeks prior to planting. Follow recommended incorporation procedures for Trilin.

PRECAUTION

Do not apply to lentils. Leaf crinkling and delayed maturity of peas may occur particularly on clay points in the northwest; but this is usually more than offset by a reduction of wild oat. Do not use foliage from treated plants for feed or grazing. Observe all precautions and limitations on the label of each product used in tank mixes and overlays.

PEANUT

Spanish Peanut in Oklahoma and Texas

Apply and incorporate Trilin prior to planting, at planting or immediately after planting using a broadcast rate of 1 pint per acre on coarse soils, and 1.5 pints on medium soils. When incorporating after planting take care not to disturb the seed.

PEPPER Transplant

Apply and incorporate Trilin prior to transplanting only.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium, 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

POTATO (Except Maine)

Apply and incorporate Trilin after planting prior to emergence or immediately following dragoff or after the potato plants have fully emerged.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium soils, 2 pints on soils with 5-10% organic matter.

Incorporation equipment should be set to uniformly cover the bed and furrow with a layer of treated soil. If the herbicide is concentrated over the bed, potato emergence may be retarded and stem brittleness can occur. Do not completely cover the foliage with treated soil when Trilin is applied and incorporated after potato plants have fully emerged. Do not completely cover foliage at subsequent cultivations. Be careful that incorporation machinery does not damage potato seed pieces or elongating sprouts.

Split Application in Idaho, Oregon and Washington

Apply and incorporate 0.75 pint of Trilin per acre before planting and 0.75 pint after planting when potato plants have fully emerged from all soils, except do not apply to soils containing a 2% or more organic matter. Follow incorporation directions listed above for application to potato after planting.

TRILIN WITH EPTAM TANK MIX in Kansas, Minnesota, Nebraska, North Dakota, Oklahoma, South Dakota and Texas

This tank mix will effectively control the following weeds in addition to those weeds controlled by Trilin.

Henbit (spring applications)	Oat, Wild
Nightshade Black	Ragweed Common
Nightshade Harry	Smartweed Pennsylvania
Nutsedge	Velvetleaf Buttonweed

Follow the recommendations for soil preparation and incorporation procedures for Trilin. Trilin with Eptam tank mix may be applied after planting but before crop emergence. In areas where potatoes are normally dragged off, this tank mix should be applied and incorporated up to or immediately following drag off.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1-1.5	1-1.5
Areas receiving greater than 20" average annual rainfall	1	1-1.5	1-2
PLUS			
Eptam 7E	1.75-7**	1.75-7	1.75-7

**For nutsedge control use the higher rate of Eptam 7E.

On soils with 2-5% organic matter use 1.5 pints per acre on coarse and medium soils and on soil with 5-10% organic matter use 2 pints on all soils.

PRECAUTION

Follow direction and precautions on the Eptam label before using. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Do not use foliage from treated plants for feed, forage or for grazing.

TRILIN WITH EPTAM APPLICATION BEFORE PLANTING in Idaho, Oregon and Washington

Trilin with Eptam may be applied prior to planting using a broadcast rate of 0.75 pint of Trilin per acre and 3.5 pints of Eptam 7E per acre on all soil textures. Incorporate immediately.

PRECAUTION

Do not use this tank mix both before and after planting in the same season. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays. Do not use foliage from treated plants for feed, forage or for grazing.

RADISH

Apply and incorporate as a preplant soil treatment.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	1.5

RAPESEED (CANOLA), CRAMBE (Except Alaska)

California - Rapeseed (Canola) Only

Follow recommended procedures for soil preparation and application of Trilin. Trilin may be applied in the fall or early spring prior to seeding. Set incorporation equipment to incorporate to a depth of 3 to 4 inches with equipment specified in this label.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2

SAFFLOWER

Recommended soil preparation application and incorporation procedures for Trilin should be followed. Use Trilin before planting in the spring or between October 15 and December 31.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

On coarse and medium soils use 1.5 pints per acre and 2 pints on fine soils with 2-5% organic matter. Use 2 to 2.5 pints on all soils with 5-10% organic matter.

For Fall Application in Arizona, Idaho, Montana, Nevada, Oregon, Utah, Washington and Wyoming

Apply and incorporate Trilin any time between October 15 and December 31. Ground may be left flat or bedded-up over winter. On bedded ground, beds should be knocked down to desired height before planting, moving some treated soil from tops into furrows. Where soil is left flat over winter, care should be taken during spring bedding operations to prevent turning up untreated soil. Destroy established weeds during seedbed preparation. Before planting destroy weeds that become established in furrows due to uncovering of untreated soil. Apply and incorporate Trilin at a broadcast rate of 1.5 pints per acre on coarse soils, 2 pints on medium and 2.5 pints on fine soils. Trilin should not be applied in the fall to soils which are wet or are subject to prolonged periods of flooding.

SOUTHERN PEA

Apply and incorporate Trilin prior to planting only.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

For soils with 2-5% organic matter use 1.5 pints on coarse and medium and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints on all soils.

SOYBEANS

Preemergence

Follow recommended soil preparation, application, and incorporation procedures for Trilin.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2

Use 1.5 pints per acre on coarse and medium textured soils and 2 pints on fine soils with 2-5% organic matter. Use 2 to 2.5 pints on all soils with 5-10% organic matter.

Fall Application

Trilin may be applied and incorporated any time between October 15 and December 31. The ground may be left flat or bedded-up over winter. Where soil is left flat, care should be taken not to turn up untreated soil from beds into furrows. On bedded grounds, beds should be knocked down to desired heights before planting, moving some treated soil into the furrow from the beds. Established weeds should be destroyed during preparation of seedbed. Before planting, destroy weeds which may have become established in furrows due to the uncovering of untreated soil during bedding. Trilin should not be applied to wet soil or soils which are subject to prolonged periods of flooding of ground where rice was grown the previous year.

Apply and incorporate Trilin at a broadcast rate of 2 pints per acre on coarse and medium soils and 2.5 pints on fine soils in Alabama, Arkansas, Northern Florida, Georgia, Louisiana, Mississippi, Southeastern Missouri, North Carolina, Oklahoma, South Carolina, Tennessee and Texas.

For other states where soybeans may be grown, use Trilin at a broadcast rate of 1 pint per acre on coarse soils, 1.5 pints on medium soils and 2 pints on fine soils. For coarse soils with 2-5% organic matter use 1.5 pints. For soils with 5-10% organic matter use 2 to 2.5 pints.

SPECIAL USE DIRECTIONS FOR SOYBEANS

Fall Panicum

Apply and incorporate Trilin broadcast at the rate of 2 pints per acre on coarse and medium soils.

Crop Rotation

Plant only rice and those crops for which Trilin can be applied as a preplant treatment following a double rate treatment season or injury may result.

Texas Panicum

Apply and incorporate at the labeled use rate for the soil type. Uniformly incorporate to a depth of at least 3 inches. Adequate soil moisture is essential for herbicide activation. Any deviation from these recommendations can result in reduced control.

Pigweed and Seedling Johnsongrass Control

In Alabama, Arkansas, Florida, Georgia, Louisiana, Mississippi, Southeastern Missouri, North Carolina, Oklahoma, South Carolina, Tennessee and Southern Virginia, apply Trilin preplant at a broadcast rate of 1 to 1.5 pints per acre on coarse soils, 1.5 to 2 pints on medium soils and 2 pints on fine soils. **Exception:** Use 3 pints per acre on fine soils in Louisiana.

Additional Weed and Grass Control

In the Texas Gulf Coast counties of Brazoria, Calhoun, Chambers, Fort Bend, Galveston, Harris, Jackson, Jefferson, Liberty, Matagorda, Orange, Victoria, Waller and Wharton, apply Trilin at a broadcast rate of 1.5 pints per acre on coarse soils, 2 pints on medium soils and 3 pints on fine soils, up to 2 weeks before planting.

Charcoal Soils in Arkansas, Louisiana and Mississippi

Freshly cleared land sometimes contains 5-10% organic matter and charcoal from burning debris. This charcoal and organic matter has a tendency to bind Trilin and reduce its weed control activity. If these conditions exist, higher rates of Trilin are needed for weed control. Crop injury can occur if increased rates are used and the charcoal or organic matter is not present in the soil. In the burn row a high level of charcoal is present, consequently, poor weed control may result even with an increased rate of Trilin.

Use Trilin broadcast at the rate of 1.5 to 2.5 pints per acre on coarse soils, 2.5 pints on medium soils and 3 pints on fine soils. Follow recommended soil preparation, application and incorporation procedures for Trilin.

Red Rice in Arkansas, Louisiana, Mississippi and Texas only

Partial suppression or control of red rice can be obtained when Trilin is applied at the following recommended rates.

Broadcast Rates Per Acre

Application	Coarse	Medium	Fine	Coarse with 2-5% organic matter	Coarse with 5-10% organic matter
Year 1	2	3	4	3	4
Year 2	1	1.5	2	1.5	2-2.5

If high organic matter and/or charcoal are present in the soil use Trilin the second year as follows for Arkansas, Louisiana and Mississippi:

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5-2.5	2.5-3

Crop Rotation

Use a 2 year program for red rice control in soybeans. Use rates listed for 1 year and plant soybeans. The second year plant only those crops for which Trilin can be used preplant using the normal rates listed for your soil type and charcoal level. Do not plant rice the second year. Rice may be planted the third year.

Rhizome Johnsongrass in Eastern United States and Texas

Rhizome johnsongrass can be acceptably controlled using a double rate program for 2 consecutive years as follows.

Soil Preparation

For satisfactory results proper soil preparation is essential. To bring rhizomes to the top of the soil use a chisel plow or similar implement. Follow twice with a disc prior to application to cut rhizomes into small 2 to 3 inch pieces and to destroy any emerged johnsongrass.

Application

Choose one of the following programs which best meets your cultural practices.

For spring application use Trilin prior to planting in the spring for 2 consecutive years. A broadcast rate of 2 pints per acre should be used on coarse soils, 3 pints on medium soils and 4 pints on fine soils. Use 3 pints on coarse soils with 2-5% organic matter and 4 pints on soils with 5-10% organic matter.

OR

For fall application use Trilin between October 15 and December 31 for 2 consecutive years using the same rates as for spring application.

OR

A split application of Trilin may be used in spring and fall for 2 consecutive years using the following rates.

Broadcast Rates Per Acre

Application	Coarse	Medium	Fine	Coarse with 2-5% organic matter	Coarse with 5-10% organic matter
Trilin (pts.)					
Spring & Fall	1	1	2	1.5	2

Incorporation

For good rhizome johnsongrass control deep incorporation is necessary. Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two passes are necessary with the second pass in a different direction from the first.

Cultivation

To control johnsongrass plants which have escaped Trilin, timely cultivations during the crop season may be necessary to obtain control. Control cannot be obtained with only 1 year of double rate Trilin use.

Crop Rotation

Plant only rice and those crops for which Trilin can be applied as a preplant treatment following a double rate treatment season or injury may result.

Wild Cane (Shattercane)

Follow recommended soil preparation and application procedures for Trilin. Germination of wild cane (shattercane) occurs throughout the growing season and from deeper in the soil than most other weed seeds. Effective control can be obtained by using the following increased rates of Trilin. A broadcast rate of 1 pint per acre on coarse soils, 2 pints on medium soils and 2.5 pints on fine soils.

Incorporation

For good wild cane control deep incorporation is necessary. Incorporate Trilin thoroughly with a disc set to cut 4 to 6 inches deep and operate at 4 to 6 mph. Two passes are necessary with the second pass in a different direction from the first.

Cultivation

Cultivation during the crop season will also contribute to control.

PRECAUTION

When using higher rates, soybeans should be planted after early season adverse weather conditions. This is to avoid additional stress to the plants due to cool, wet weather early in the growth period. Use of high rates and adverse growing conditions can cause reduced stands, delayed maturity and reduced yields.

TANK MIXES AND OVERLAYS

Observe all precautions and limitations on the labels of each product used in tank mixes and overlays.

TRILIN WITH SENCOR OR LEXONE

Trilin with Sencor or Lexone will control certain grasses and broadleaf weeds listed for Trilin alone and the following weeds:

Jimsonweed	Ragweed, Common
Mallow, Venice	Sesbania, Hemp
(Flower-of-an-hour)	Smartweed, Pennsylvania
Mustard, Wild	Velvetleaf
Prickly Sida	

Cocklebur, morningglory and giant ragweed (horseweed) control may be erratic. Timely cultivation may improve control. An overlay of Sencor or Lexone may be preferred to the tank mix where cocklebur is a serious problem.

Trilin tank mixed with Sencor or Lexone may be applied from 2 weeks prior to planting up to planting.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2
PLUS			
Sencor 4L (pts.)	0.5	0.75	1
OR			
Lexone DF or			
Sencor DF (lb.)	0.33	0.5	0.67

Do not use Sencor or Lexone on coarse soils with less than 1% organic matter.

PRECAUTION

Do not plant any crop other than soybeans within 4 months after treatment. Crop injury, herbicide residue or erratic weed control may result from over application, uneven application or improper soil incorporation. Cold weather, deep planting, seedling disease, excessive moisture, soil pH over 7.5, high salt concentration or drought are additional stress factors. Any of these may weaken crop seedlings and possible damage from the tank mix is increased. These factors may also delay crop development or reduce yields when Sencor or Lexone is applied. Do not use foliage from treated plants for feed on forage.

TRILIN PREPLANT FOLLOWED BY SENCOR OR LEXONE AS AN OVERLAY

After Trilin has been applied as a preplant incorporated herbicide make a single application of Sencor or Lexone as either a broadcast or band spray either during planting or after planting, but before the soybeans emergence. Crop injury may result if Sencor or Lexone is sprayed over the top of emerged soybeans.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)	1	1.5	2
PLUS			
Sencor 4L (pts.)	0.75-1	0.75-1.5	1-1.75
OR			
Sencor DF (lbs.)	0.5-0.67	0.5-1	0.67-1.17
OR			
Lexone DF (lb.)	0.5	0.5-0.67	0.67

Lexone or Sencor should not be applied to sands or soils with less than ½% organic matter or to coarse soils (sandy loam and loamy sand) containing less than 2% organic matter.

PRECAUTION

Tracy, Semmes, Altona, Vansoy or Coker 102 soybean varieties are sensitive and crop injury may result if Lexone or Sencor is used. Before a Sencor or Lexone application, seeds must be planted at least 1.5 inches but not more than 2 inches below the soil surface. Only one application per season should be used at these rates. Do not plant areas treated with Sencor or Lexone to any crop other than soybeans within 4 months after treatment. Soybean injury may occur if Lexone or Sencor is used on soils having a calcareous surface, a pH of 7.5 or higher, or if they are used in conjunction with soil-applied organic phosphate pesticides. Do not use foliage from treated plants for feed or forage. Observe all precautions and limitations on the labels of each product used in tank mixes and overlays.

SUGARBEET

Use Trilin when plants are between 2 to 6 inches tall as a broadcast over-the-top spray at a rate of 1 pint per acre on coarse soils and 1.25 to 1.5 pints on medium and fine soils. Use the higher rate for medium and fine soils in areas receiving greater than 20 inches average annual rainfall. Set incorporation machinery to throw treated soil toward the plants in the row. Care should be taken that incorporation machinery does not damage the sugarbeet taproot.

PRECAUTION

To reduce the possibility of girdling, exposed beet roots should be covered with soil before applying Trilin.

In Colorado, Idaho, Montana, Nebraska, Oregon, Texas, Utah, Washington and Wyoming use a tine-tooth harrow (Flexline or Melroe) for incorporation for Trilin for effective weed control in sugarbeets. The tine-tooth harrow should be operated two times over the field, the second time in opposite direction at a speed of 3 to 6 mph. The harrow should be set to cut 1 or 2 inches deep. Care should be taken that the tine-tooth harrow does not damage the sugarbeet taproot. All recommendations for application procedures and broadcast rates per acre for sugarbeet should be followed.

SUGARCANE Plant Cane

Apply and incorporate Trilin twice a year at a broadcast rate of 2 to 4 pints per acre for all soil textures. Make the first application of Trilin in the fall immediately after the seed pieces are planted and the second application of Trilin in the spring before or shortly after the cane emerges. Before the spring application rain-packed beds should be loosened 2 to 3 inches deep. Care should be taken so that the seed pieces or emerging shoots are not damaged by incorporation machinery.

Hawaii (postplant) for control of most annual grasses including guineagrass

Surface apply Trilin after planting (for plant cane) or after harvesting (for ratoon cane) at a broadcast rate of 6 to 8 pints per acre for all soil textures. Apply Trilin prior to germination and emergence of grass weeds. For best results in plant cane, the soil surface should be smooth and finely tilled. Apply Trilin as soon as possible after tillage and planting, before germination and emergence of grass weeds. For optimum efficacy in ratoon cane, minimize surface residue from previous crop before applying. For best result, apply Trilin before anticipated rainfall in non-irrigated and furrow-irrigated sugarcane. Apply 0.5 inch or more irrigation in drip-irrigated or sprinkler-irrigated sugarcane as soon as possible after applying Trilin.

Repeat Applications

Subsequent germination flushes of grass weeds may occur prior to the development of a full dense canopy of sugarcane. Once this occurs, additional grass weed establishments are strongly suppressed. One or two additional applications of Trilin can be applied to maintain weed control during the early crop development period. For repeat applications, direct the spray to the soil surface to minimize interception of the herbicide by the crop.

PRECAUTION

Do not apply Trilin as a postplant surface applied treatment within 180 days of harvest.

Applications in Louisiana or Texas up to lay-by for plant cane or ratoon cane

Use Trilin at a broadcast rate of 2 to 4 pints per acre for all soil textures. This should be done in the spring from before or shortly after the cane emerges up to lay-by. Use Trilin after the beds have been shaved or false shaved. Rain-packed beds should be loosened 2 to 3 inches deep before application. Care should be taken that seed pieces or emerging shoots are not damaged by incorporation machinery. A rolling cultivator or bed chopper may be used to incorporate Trilin lay-by applications in sugarcane on all soil textures. Normal incorporation directions for the rolling cultivator should be followed. Bed chopper should be set to cut 3 to 4 inches deep and operate at 4 to 6 mph. Two incorporation passes are necessary.

Itchgrass (Raoulgrass) control in Louisiana

Apply and incorporate Trilin on either plant or ratoon cane at a broadcast rate of 4 pints per acre for all soil textures. Directions above for sugarcane lay-by application in Louisiana and Texas should be followed.

SUNFLOWER

Use Trilin in the spring or in the fall between October 15 and December 31. Recommendations for soil preparation application and incorporation procedures for Trilin should be followed.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1.5
Areas receiving greater than 20" average annual rainfall	1	1.5	2

On coarse and medium soils use 1.5 to 2 pints and 2 pints on fine soils with 2-5% organic matter. Use 2 pints on all soils with 5-10% organic matter.

TOMATO

For direct-seeded tomato use Trilin at blocking or thinning as a directed spray to the soil between rows and beneath the plants and incorporate. For transplant tomato, apply and incorporate Trilin preplant, postplant or at lay-by.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Areas receiving less than 20" average annual rainfall	1	1.25-1.5	1-2
Areas receiving greater than 20" average annual rainfall	1	1.5	2

On soils with 2-5% organic matter use 1.5 pints on coarse and medium textured soils and 2 pints on fine soils. On soils with 5-10% organic matter use 2 pints.

VINEYARDS

See FRUIT AND NUT CROPS AND VINEYARDS for instructions.

WHEAT (WINTER)

Colorado, Idaho, Kansas, Montana, Nebraska, Oregon, Washington and Wyoming

Apply Trilin for preplant preemergence control of cheatgrass and other annual grasses and broadleaf weeds controlled by Trilin. If the seed is placed below the zone of soil treated with Trilin the growth development and yield of winter wheat will not be adversely affected.

Use Trilin any time during a period from 3 weeks up to immediately prior to planting. Apply Trilin at a broadcast rate of 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils.

Incorporation

Incorporate Trilin into the soil with a flexible tine-tooth harrow (Flexline or Melroe) set to cut 1 to 2 inches deep and operate at 3 to 6 mph. Within 24 hours after application incorporate one time followed by a second incorporation in a different direction from the first prior to planting. After the Trilin has been incorporated with a flexible tine harrow do not till the soil with a disc.

Seeding Directions

Use only a deep furrow or semi-deep furrow drill that will place the seed below the zone of soil into which Trilin has been incorporated.

PRECAUTION

Crop injury such as delayed emergence and development may occur when wheat is planted in direct contact with treated soil.

Fallow soil application in Washington and Oregon

To control cheatgrass and certain annual grasses and broadleaf weeds, apply Trilin and shallowly incorporate into fallow soil up to 4 months ahead of planting. As long as the seed is placed below the zone of soil treated with Trilin the growth development or yield will not be adversely affected. Use deep or semi-deep furrow drills. Use at a broadcast rate of 1.5 pints per acre on coarse and medium soils and 2 pints on fine soils. Trilin can be applied any time from May to September prior to the fall planting of winter wheat.

Incorporation

Incorporate Trilin using a flexible tine-tooth harrow (Flexline or Melroe) set to cut 1 to 2 inches deep and operated at 3 to 6 mph. Two passes over the field in different directions are necessary for thorough incorporation. Incorporate one time within 24 hours after application followed by a second incorporation prior to seeding. When a flexible tine harrow has been used to apply Trilin do not till the soil with a disc.

PRECAUTION

Deep furrow or semi-deep furrow drills only should be used. Seed should be placed below the zone of soil into which Trilin has been incorporated. Injury to the crop or delay in emergence or development may occur if wheat is planted directly into the zone of soil treated with Trilin.

WHEAT (SPRING), DURUM AND BARLEY

Preplant Barley

Apply and incorporate Trilin before planting in the spring or in the fall after September 1, at the rate of 1 pint per acre on coarse soils, 1.5 pints per acre on medium soils and 1.5 pints on fine soils.

Use 1.5 pints of Trilin per acre on coarse and medium soils with 2-5% organic matter. Incorporate Trilin into the soil with a flexible tine-tooth harrow, (Flexline or Melroe) set to cut 1 to 2 inches deep and operate at 3 to 6 mph. Within 24 hours after application, incorporate one time followed by a second incorporation in a different direction from the first, prior to planting. After the Trilin has been incorporated with a flexible tine harrow do not till the soil with a disc. Use only a deep furrow or semi-deep furrow drill that will place the seed below the zone of soil into which Trilin has been incorporated.

PRECAUTION

Crop injury such as delayed emergence and development may occur when barley is planted in direct contact with treated soil. DO NOT MAKE MORE THAN ONE TRILIN APPLICATION PER SEASON.

Postemergence Wheat (Spring), Durum and Barley

Trilin Alone

To control foxtail (pigeongrass) Trilin is recommended as a postplant incorporated treatment. Use Trilin at a broadcast rate of 1 pint per acre on coarse and medium soils and 1.5 pints on fine soils. Seedbed should be well-tilled and seed planted 2 to 3 inches deep. Use Trilin after seeding but prior to crop emergence. Use flexline or diamond harrow operated two times in different directions to incorporate. Incorporate by operating equipment at a speed of at least 5 mph and set at up to 1.5 inches deep. Apply and incorporate the first time in the same operation if possible. Both incorporations must be done within 24 hours.

TRILIN WITH FAR-GO TANK MIX

This tank mix will effectively control all the weeds controlled by Trilin alone plus these additional weeds:

Foxtail	Pigeongrass	Wild Oat
---------	-------------	----------

Apply Trilin with Far-Go as a postplant incorporated treatment. Plant 2 to 3 inches deep in a well-tilled seedbed. Trilin with Far-Go should be applied after seeding but prior to crop emergence. Use flexline or diamond harrows to incorporate. Make two passes each in different directions at speeds of at least 5 mph operating equipment 1 to 1.5 inches deep. Application and the first incorporation should be done in the same operation if possible. If not incorporate immediately after application.

Broadcast Rates Per Acre

	Coarse	Medium	Fine
Trilin (pts.)			
Barley, Durum, Spring Wheat PLUS Far-Go (pts.)			
Durum, Spring Wheat	2.5	2.5	2.5
Barley	2	2	2

PRECAUTION

Over application may result in crop injury. Observe all precautions and limitations on the label of each product used in tank mixes and overlays. DO NOT MAKE MORE THAN ONE TRILIN APPLICATION PER SEASON.

FERTILIZER USE DIRECTIONS

LIQUID FERTILIZERS

Trilin may be mixed with most liquid fertilizer materials. A combination of Trilin with solutions and suspension-type fertilizers will provide weed and grass control equal to the same rates of Trilin applied to water. Trilin label recommendations regarding rates per acre, crops, incorporation directions, special instructions, cautions and special precautions should be followed. Labeling and applications relating to liquid fertilizer mixing for individual state regulations are the responsibility of the individual and/or company selling the fertilizer and chemical mixture.

Testing for Tank Mix Compatibility In Liquid Fertilizers

A tank mixture of Trilin alone or with dry flowables, wettable powders, aqueous suspensions, flowables liquids or solutions may not combine properly with some liquid fertilizer materials. Always test a small quantity before full-scale mixing to determine whether a compatibility agent is needed and which agent does the best job. Phosphate ester type surfactants designed for use with liquid fertilizers are suggested. Use the following test to select the correct agent for your mixture.

1. Measure 1 pint of intended spray water or fertilizer solution into a jar.
2. Check the pH of the liquid and adjust if necessary.
3. Add in the given order the intended ingredients shaking well after each addition.
 - (a) surfactants (spreaders), acidifies, compatibility agents and activators; add 1 teaspoon for each pint per 100 gallons.
 - (b) dry ingredients (wetable powders or dry flowables); add 1 tablespoon for each pound per 100 gallons.
 - (c) emulsifiable concentrates; add 1 teaspoon for each pint per 100 gallons.

Rate Chart For Impregnating Fertilizer With Trilin Added to a Ton of Fertilizer

CHART QUANTITIES LISTED ARE QUARTS OF TRILIN PER TON OF FERTILIZER

Fertilizer Rate Per Acre	TRILIN Rate Per Acre				
	1 pt.	1½ pts.	2 pts.	3 pts.	4 pts.
200 lbs.	5	7.5	10	15	20
250 lbs.	4	6	8	12	16
300 lbs.	3.33	5	6.67	10	13.33
350 lbs.	2.75	4.25	5.75	8.5	11.5
400 lbs.	2.5	3.75	5	7.5	10
450 lbs.	2.25	3.33	4.5	6.67	9

For rates other than those listed above use the following formula to calculate the amount of Trilin to be impregnated on a ton of dry bulk fertilizer

$$\begin{array}{rclclcl} \text{Pts.} & & & & & \text{Qts.} \\ \text{Trilin/Acre} & \times & 1000 \text{ lbs.} & = & \text{Trilin/Ton of} & \\ & & \text{Fertilizer/Acre} & & \text{Fertilizer} & \end{array}$$

All individual state regulations relating to dry bulk fertilizer blending, registration, labeling and application are the responsibility of the individual and/or company selling the fertilizer and chemical mixture.

Application

Use a properly calibrated applicator to spread the fertilizer/chemical mixture. The material should be applied uniformly to the soil surface.

Incorporation

Follow Trilin incorporation procedures.

NOTICE OF WARRANTY

GRIFFIN warrants that this product conforms to the chemical description on the label thereof and is reasonably fit for purposes stated on such label only when used in accordance with directions under normal use conditions. It is impossible to eliminate all risks inherently associated with use of this product. Crop injury, ineffectiveness or other unintended consequences may result because of such factors as weather conditions, presence of other materials or the manner of use or application, all of which are beyond the control of GRIFFIN. In no case shall GRIFFIN be liable for consequential, special or indirect damages resulting from the use or handling of this product. All such risks shall be assumed by the Buyer. The exclusive remedy of any buyer or user of this product for any and all losses, injuries, or damages resulting from or in any way arising from the use, handling, or application of this product, whether in contract, warranty, tort, negligence, strict liability, or otherwise, shall not exceed the purchase price paid for this product or at GRIFFIN'S election, the replacement of this product. GRIFFIN MAKES NO WARRANTIES OF MERCHANTABILITY OR FITNESS FOR A PARTICULAR PURPOSE NOR ANY OTHER EXPRESS OR IMPLIED WARRANTY EXCEPT AS STATED ABOVE.

Caparol is a registered trademark of Ciba-Geigy Corporation. GCN 102998
Cotoran is a registered trademark of Ciba-Geigy Corporation. CPC 026563
Cotton-Pro is a registered trademark of Griffin Corporation.
Direx 4L is a registered trademark of Griffin Corporation.
Eptam is a registered trademark of ICI Americas, Incorporated.
Far-Go is a registered trademark of Monsanto Agricultural Products Company.
Griffin and Design are a registered trademark of Griffin Corporation.
Karmex is a registered trademark of E.I. Dupont de Nemours and Company.
Lexone is a registered trademark of E.I. Dupont de Nemours and Company.
Meturon is a registered trademark of Griffin Corporation.
Sencor is a registered trademark of Mobay Corporation.
Trilin is a registered trademark of Griffin Corporation.

- (d) flowables: add 1 teaspoon for each pint per 100 gallons.
 - (e) soluble ingredients: add 1 tablespoon for each pound per 100 gallons.
 - (f) spreader-stickers: 1 teaspoon for each pint per 100 gallons.
4. The final mixture should be uniform and smooth with no evidence of coagulation occurring. If incompatibility is evident begin test again with a compatibility agent added first. Six drops is equivalent to 4 ounces per 100 gallons. If this does not smooth the mixture try higher concentrations and other compatibility agents.
5. Allow the mixture to stand undisturbed 30 minutes. If separation occurs shake and observe the resulting mixture. If the mixture does not redisperse do not attempt to spray the mixture. You may try:
- (a) more compatibility agents
 - (b) different formulations of the active ingredients (switch from wettable or emulsifiable concentrates to flowable or from wettable powder to emulsifiable concentrates).

Mixing Instructions

When mixing Trilin in liquid fertilizers continuous agitation is required to prevent the Trilin from rising to the surface as an oily layer. Use a compatibility agent to make the Trilin emulsify properly.

When tank mixing emulsifiable concentrates with dry flowables, wettable powders, aqueous suspensions, flowables liquids or solutions in liquid fertilizer using a compatibility agent is especially important. If Trilin rises to the surface of the fertilizer as an oil and the emulsion is not properly formed, the oil may combine with the formulation or suspension to form oil curds which is difficult to redisperse. A compatibility agent is helpful in causing liquid concentrates to form non-oiling mixtures with liquid fertilizers. Compatibility agents can be used at rates as low as 1.5 to 2 pints per ton of liquid fertilizer and should be mixed well with the fertilizer before adding the liquid concentrate. Follow the directions on the compatibility agent label.

The following is a list of some phosphate ester type surfactants designed to be used with liquid fertilizers. They usually do not work as compatibility agents in tank mixtures in plain water.

*Amoco Spray Mate (Amoco Oil Company, Chicago, IL)

*Compat (Farm Chemicals Incorporated, Aberdeen, NC)

Kem-Link (Universal Coop., Minneapolis, MN)

Rigo Compatibility Agent (Rigo Company, Buckner, KY)

Sponto 168D (Witco Chemicals Company, Chicago, IL)

*T-Mulz 734-2 (Thompson-Hayward Chemical Co.)

Unite (Hopkins Ag Chemicals, Madison, WI)

*Not for use in California

Application

Use a properly calibrated applicator to spread the fertilizer/pesticide mixture and apply material uniformly to the soil surface.

Incorporation

Follow normal Trilin incorporation procedures.

DRY BULK FERTILIZERS

Trilin may be used for impregnation or coating of dry bulk fertilizers. Application of dry bulk fertilizers impregnated with Trilin has provided weed and grass control equal to the same rates of Trilin applied in water. Follow all Trilin label recommendations regarding rates per acre, approved crops, incorporation, special instructions, cautions and special precautions. A minimum of 200 pounds per acre of dry fertilizer impregnated with Trilin at the recommended rates should be applied. Trilin can be used for impregnation of any commonly used fertilizer, except coated ammonium nitrate and straight limestone. Trilin will not be absorbed by these materials. Blends containing mixtures of these materials can be impregnated.

Impregnation

A closed drum belt ribbon or other commonly used dry bulk fertilizer blender may be used. Trilin should be applied uniformly to the fertilizer.

Rates

Specific crop recommendations for the rate of Trilin per acre should be followed. Check the rate table below to determine the amount of Trilin to be impregnated into a ton of dry bulk fertilizer based on the amount of fertilizer which will be applied per acre.