

[Translated by Julian Brooke and published with the permission of the Chinese Journal of Epidemiology 1994 October;15(4):296-8.]

THE EFFECTS OF HIGH LEVELS OF FLUORIDE AND IODINE ON CHILD INTELLECTUAL ABILITY AND THE METABOLISM OF FLUORIDE AND IODINE

Yingkui Yang, Xiuhong Wang, Xiaowei Guo, Peiying Hu^a
Jinan, China

SUMMARY: The authors carried out a study on the intellectual abilities and fluoride/iodine metabolism of children living in a high fluoride-high iodine area. Among the results: the percentage of the general population living in this fluoride/iodine-contaminated region that suffered from goiter (clinical thyroid enlargement) was 3.8%, the rate of children already showing some thyroid enlargement was 29.80%. Similarly, the rate of dental fluorosis for the general population was 35.48%, while for children it was 72.98%. Student subjects had average IQs of 76.67 ± 7.75 , with 16.67% of the IQs in the "low" category. The iodine content and fluoride content of the children's urine were $816.25 \pm 1.80 \mu\text{g/L}$ and $2.08 \pm 1.08 \text{ mg/L}$, respectively, significantly higher than the control. The thyroid glands of the subjects showed a markedly lower uptake rate of iodine-131 when compared with the control, the values after 3 hr and 24 hr were $9.36 \pm 1.55\%$ and $9.26 \pm 4.63\%$, respectively, and the blood serum levels of thyroid stimulating hormone (TSH) were significantly higher than the control. The results indicate that high levels of fluoride and iodine have a serious damaging effect on the body, and should be given greater attention.

Keywords: Child IQ; Dental Fluorosis; High fluoride water; High iodine intake; Iodide Goiter; Thyroid Stimulating Hormone.

INTRODUCTION

The subjects of this study were residents of the Huimin and Dezhou regions of Shandong Province located on the lower reaches of the Yellow River. In 1976, in a response to increasingly bitter, salty water, much deeper wells were dug. However, the fluoride and iodine content of this deep well water was found to be much higher than the standards for drinking water, resulting in a high prevalence of iodine goiter and fluoride poisoning. The area is known locally as the "twin contamination zone." In order to investigate the effects of high iodine and fluoride on child intellectual ability and the metabolic characteristics of iodine and fluoride, we carried out a comparative study of two villages in Qingyun County with the following results.

MATERIALS AND METHODS

1. Lidian village of Qingyun County was selected as the site of our study; its drinking water was tested to have an iodine content of $1,100 \mu\text{g/L}$ and a fluoride content of 2.97 mg/L . The non-disease control was Dading village of the same county; its drinking water showed an iodine content of $128.6 \mu\text{g/L}$ and a fluoride content of 0.5 mg/L .

2. The diagnoses of goiter and dental fluorosis were both carried out according to national standards for endemic disease control.^{1,2} Intelligence testing was done using the Chinese Comparative Scale of Intelligence Test (Third Edition), as revised by Wu Tianming.³

3. The determination of iodine and fluoride levels in the drinking water was accomplished by conventional physicochemical analysis techniques, with their values expressed in units of $\mu\text{g/L}$ and mg/L , respectively. The incineration method was used to determine urinary iodine, again expressed as $\mu\text{g/L}$, while the electrode

^aThe authors work at the Shandong Provincial Institute of Endemic Disease Control, Jinan, 250014, PR China.

which was somewhat lower than the control (IQ 81.67 ± 11.9), although the difference is not statistically significant ($P > 0.05$). However, as seen in Table 2, the percentage of children in the low range (16.67%) is higher in the endemic group than in the control group (10.0%), suggesting that a high iodine-high fluoride environment also has a definite negative influence on child intellectual ability.^c

The urinary iodine and urinary fluoride levels for children living in the high iodine-high fluoride area were $816.25 \pm 1.80 \mu\text{g/L}$ and $2.08 \pm 1.03 \text{ mg/L}$, respectively, clearly elevated by comparison to the control, reflecting the high body load of iodine and fluoride. Elemental iodine is a key component of thyroid hormones, and also influences several stages of their formation and excretion.⁵ Fluoride, on the other hand, is toxic to living cells, and is a powerful inhibitor of certain enzymes. Excessive uptake of fluoride can cause decreased functioning of the thyroid gland in direct relation to the blood level. Of course, iodine is the primary factor here; what exact role that fluoride is playing requires further study.

The serum levels of T_3 and T_4 for the children from the high iodine-high fluoride zone were only slightly higher than the control ($P > 0.05$), but the level of TSH was clearly elevated ($P < 0.01$). That the thyroid gland excretion of T_3 and T_4 for the "twin contamination zone" children was in the normal range, but the pituitary gland secretion of TSH was significantly elevated making it probable that reverse feedback is promoting the hypothalamus excretion of TRH (thyroid releasing hormone), causing a corresponding increase in the excretion of TSH, which stimulates compensating production of T_3 and T_4 by the thyroid gland.

In summary, the results of this study indicate the following:

(1) Areas that have long-term, serious iodine and fluoride contamination can cause goiter and fluoride poisoning in the population; this problem should be taken seriously.

(2) The clinical characteristics of children from this region include high urinary iodine, high urinary fluoride, poor thyroid iodine-131 absorption with similar values at 3 and 24 hr (possibly a backward peak value shift), and high TSH values.

(3) With regard to the diseases endemic to this high iodine-high fluoride zone, the only fundamental means of control is to change the source of water, limit the ingestion and absorption of iodine and fluorine, and promote their excretion from the body.

REFERENCES

- 1 The North China Professional Council on Iodized Salt as a Means of Controlling Endemic Goiter. Standards for work to control endemic goiter (Trial Version). Chinese Journal of Preventive Medicine 1978;12(1):62.
- 2 Dai Guojun. Endemic fluoride poisoning. Hohot: The Inner Mongolian People's Publishing House; 1985. p. 383-7.
- 3 Wu Tianmin. The Chinese Comparative Intelligence Test Guidebook (3rd revision). Peking University Press; 1983. p. 1-91.
- 4 Ren Dali, Li Kecheng, Liu Dawei. Study on the intellectual development of 8-14 year-old children in high fluoride, low iodine zones. Chinese Journal of Control of Endemic Diseases 1989;4(4):351. [A translation of this paper is in this issue of Fluoride on p. 319-20.]
- 5 Yang Yingkui, Zhang Xining, Zhang Huifang, et al. Research on iodine metabolism, pituitary/thyroid function, and the pathology of goiter in patients exposed to excess iodine. Chinese Journal of Control of Endemic Diseases 1988;3(6):338.
- 6 Liu Zhenqi, Hu Min, Wu Hanwen, et al. Investigation of the relationship between serum thyroid hormone and fluoride levels in fluoride-poisoned patients. Chinese Journal of Control of Endemic Diseases 1988;7(4):216.

^cAn editorial comment on the statistical testing is given as a footnote on p. 337.